

Topological regluing of rational functions

V. Timorin

Received: 17 October 2008 / Accepted: 12 October 2009 /
Published online: 10 November 2009
© Springer-Verlag 2009

Abstract Regluing is a surgery that helps to build topological models for rational functions. It also has a holomorphic interpretation, with the flavor of infinite dimensional Thurston–Teichmüller theory. We will discuss a topological theory of regluing, and just trace a direction, in which a holomorphic theory can develop.

1 Introduction

1.1 Overview and main results

Consider a continuous map $f : S^2 \rightarrow S^2$. We are mostly interested in the case, where f is a rational function of one complex variable considered as a self-map of the Riemann sphere. The objective is to study the topological dynamics of f . In particular, how to modify the topological dynamics in a controllable way? There is an operation that does not change the dynamics at all: a conjugation by a homeomorphism. Let $\Phi : S^2 \rightarrow S^2$ be a homeomorphism, and consider the map $g = \Phi \circ f \circ \Phi^{-1}$. Then one can think of g as being “the map f but in a different coordinate system”. In particular, all dynamical properties of the two maps are the same. Another example is a semi-conjugacy. Let $\Phi : S^2 \rightarrow S^2$ now be a continuous surjective map, but

V. Timorin (✉)
Faculty of Mathematics, Higher School of Economics, 20 Myasnitskaia St., 101000
Moscow, Russia
e-mail: vtimorin@hotmail.com

V. Timorin
Jacobs University Bremen, 28759 Bremen, Germany

not a homeomorphism. Sometimes, the “conjugation” $g = \Phi \circ f \circ \Phi^{-1}$ still makes sense, although Φ^{-1} does not make sense as a map. Namely, this happens when f maps fibers of Φ to fibers of Φ . In this case, g is well-defined as a continuous map. Such map is said to be *semi-conjugate* to f .

If we want to perform a surgery on f , then we need to consider discontinuous maps Φ (*if you don't cut, that is not a surgery*). Of course, if we allow badly discontinuous maps Φ , then the “conjugation” $g = \Phi \circ f \circ \Phi^{-1}$, even if it makes sense as a map, would have almost nothing in common with f , in particular, it would be hard to say anything about the dynamics of g . Thus we must confine ourselves with only nice types of discontinuities. An example is the following: given a simple curve, one can cut along this curve, and then reglue in a different way.

E.g. consider the following map:

$$j(z) = \sqrt{z^2 - 1}.$$

Note that there are two branches of this map that are well-defined and holomorphic on the complement to the interval $[-1, 1]$. We choose the branch that is asymptotic to the identity near infinity and call it j . The map j has a continuous extension to each “side” of the interval $[-1, 1]$ but the limit values at different sides do not match. By considering the limit values of j at both sides of $[-1, 1]$, we can say that j reglues this interval into the interval $[-i, i]$ (i.e. the straight line segment connecting i and $-i$). A precise definition of a regluing will be given in Sect. 2.1. For now, a regluing of a set of disjoint simple curves is a one-to-one map defined and continuous on the complement to these curves and behaving near each curve as the map j considered above.

We are forced to consider regluings of countably many curves. In fact, if we want to reglue some curve in the dynamical picture of a function f , then, in order to have a global continuous extension of $\Phi \circ f \circ \Phi^{-1}$, our regluing Φ must also reglue all pullbacks of this curve under f .

Below, we briefly explain the main result. Since the precise definitions are rather lengthy, we will only give a sketch, postponing the detailed statements until the main body of the paper. Let Φ be a regluing of countably many disjoint simple curves in the sphere (note that the complement to countably many simple curves is not necessarily open but is always dense). Also, consider a branched covering $f : S^2 \rightarrow S^2$. Under certain simple topological conditions on f and the curves, we can guarantee that the map $\Phi \circ f \circ \Phi^{-1}$ extends to the whole sphere as a branched covering. We say that this covering is obtained from f by *topological regluing of disjoint simple curves*. With the help of topological regluing, new topological models of rational functions can be obtained from the existent models.

The simplest example is provided by quadratic polynomials. E.g. consider the quadratic polynomial $f(z) = z^2 - 6$ (the particular choice of number 6

does not have any importance; we could as well take any real number bigger than 2). Most points escape to infinity under the iterations of f . The set of points that do not escape is a Cantor set J_f called the *Julia set of f* . This set lies on the real line. The right-most point of J_f is 3. Note that 3 is fixed under f . The left-most point of J_f is -3 , which is mapped to 3. The biggest component of the complement to J_f in $[-3, 3]$ is $(-\sqrt{3}, \sqrt{3})$, and all other components are pullbacks of $(-\sqrt{3}, \sqrt{3})$ under the iterates of f . The points $\pm\sqrt{3}$ are mapped to -3 . Suppose that we reglue the interval $[-\sqrt{3}, \sqrt{3}]$ and all its pullbacks under f . Then the Julia set of f collapses into a connected set homeomorphic to the interval. Moreover, we can choose the corresponding regluing map Φ in such a way that $\Phi \circ f \circ \Phi^{-1}$ extends to the quadratic polynomial $g(z) = z^2 - 2$, the so called *Chebyshev polynomial*. The Julia set of g is the interval $[-2, 2] = \Phi(J_f)$. We will work out this example in detail in Sect. 5.1.

More generally, let f be a quadratic polynomial $z \mapsto z^2 + c$, where c is the landing point of an external parameter ray \mathcal{R} . Suppose that the Julia set of f is locally connected, and all periodic points of f in \mathbb{C} are repelling. Also, consider a quadratic polynomial g , for which the corresponding parameter value belongs to \mathcal{R} . Thus the Julia set of g is disconnected. Then f and g can be obtained one from the other by a regluing (of a set of disjoint simple curves). This is explained in Sect. 2.3.

However, the main motivation for the notion of regluing was the problem of finding topological models for quadratic rational functions. According to a well-known general observation, the dynamical behavior of a rational function is determined by the behavior of its critical orbits. A quadratic rational function has two critical points. Thus, to simplify the problem, one puts restrictions on the dynamics of one critical point, and leaves the other critical point “free”. For example, it makes sense to consider quadratic rational functions with one critical point periodic of period k . For $k = 1$, we obtain quadratic polynomials. Indeed, the fixed critical point can be mapped to infinity, and a quadratic rational function having infinity as a fixed critical point is necessarily a quadratic polynomial.

Suppose now that $k > 1$, and f is a quadratic rational function with a k -periodic critical point and the other critical point non-periodic. Recall that f is called a *hyperbolic rational function of type B* if the non-periodic critical point of f lies in the immediate basin of the periodic critical cycle (but necessarily not in the same component, see e.g. [8, 12]). The function f is said to be a *hyperbolic rational function of type C* if the non-periodic critical point of f lies in the full basin of the periodic critical cycle, but not in the immediate basin. The classification of hyperbolic rational functions into types was introduced by M. Rees [12]. However, a different terminology was used (types II and III instead of types B and C). We use the terminology of

Milnor [8], which is more popular and perhaps more suggestive (B stands for “Bi-transitive”, and C for “Capture”).

Fix $k > 1$. The set of hyperbolic rational functions with a k -periodic critical point splits into *hyperbolic components*. We say that a hyperbolic component is of type B or C if it consists of hyperbolic rational functions of this type. There are also type D components, which we do not discuss in this paper. There are no type A components for $k > 1$.

Theorem 1 *Let f be a quadratic rational function with a k -periodic critical point. If f is on the boundary of a type C hyperbolic component but not on the boundary of a type B hyperbolic component, then f is the continuous extension of $\Phi \circ h \circ \Phi^{-1}$ to the sphere, where h is a critically finite hyperbolic rational function, and Φ is a regluing of a countable set of disjoint simple curves. Moreover, h can be chosen to be the center of any type C hyperbolic component, whose boundary contains f .*

This result, combined with the topological models for hyperbolic critically finite functions given in [13], provides topological models for most functions on the boundaries of type C components. We will prove Theorem 1 in Sect. 3. The requirement that f be not on the boundary of a type B component is probably inessential. However, to study functions on the boundary of a type B component, one needs to use different techniques. For $k = 2$, there is only one type B component, and a complete description of its boundary is available [17]: all functions on the boundary are simultaneously matings and anti-matings. See also [1, 5] for other interesting results concerning the case $k = 2$. On the other hand, I do not know any example of a type C hyperbolic component and a type B hyperbolic component, whose boundaries intersect at more than one point. In the proof of Theorem 1, many important ideas of [13] are used. At some point, we employ an analytic continuation argument similar to that in [1].

In Sect. 6, we prove that under some natural assumptions on a set of simple disjoint curves on the sphere, there exists a topological regluing of this set. This statement is a major ingredient in the proof of Theorem 1. The existence result is based on a theory of Moore [10], which gives a topological characterization of spaces homeomorphic to the 2-sphere.

In Sect. 5, we define an explicit sequence of approximations to a regluing. These approximations are defined and holomorphic on the complements to finitely many simple curves (not necessarily disjoint). The holomorphy of approximations may prove to be important. However, we just introduce the basic notions and postpone a deeper theory for future publications.

2 Topological regluing

We define topological regluing and give simplest examples, in particular, we discuss topological regluing of quadratic polynomials.

2.1 Definition and examples

Let S^1 denote the unit circle in the plane. In Cartesian coordinates (t_1, t_2) , it is given by the equation $t_1^2 + t_2^2 = 1$. We will also consider the sphere S^2 obtained as the one-point compactification of the (t_1, t_2) -plane. In this sphere, consider the region Δ_∞ bounded by S^1 and containing ∞ (i.e. the outside of the unit circle). The closure of this region in the sphere is denoted by $\overline{\Delta}_\infty$. We will write Δ for the unit disk $\{t_1^2 + t_2^2 < 1\}$. The Klein 4-group V_4 acts on the unit circle. Namely, the two generators of V_4 are $s_1 : (t_1, t_2) \mapsto (t_1, -t_2)$ and $s_2 : (t_1, t_2) \mapsto (-t_1, t_2)$.

A continuous map $\alpha : S^1 \rightarrow S^2$ is called an α -path if $\alpha(t_1, t_2) = \alpha(t'_1, t'_2)$ is equivalent to $t'_1 = t_1$. Note that $\alpha(t_1, t_2)$ is then an injective continuous function of t_1 , and that $t'_1 = t_1$ implies $t'_2 = \pm t_2$. Similarly, a continuous map $\beta : S^1 \rightarrow S^2$ is called a β -path if $\beta(t_1, t_2) = \beta(t'_1, t'_2)$ is equivalent to $t'_2 = t_2$. Thus β can be represented as an injective continuous function of t_2 . Note that every simple path $\gamma : [-1, 1] \rightarrow S^2$ can be interpreted either as an α -path defined as $\alpha(t_1, t_2) = \gamma(t_1)$ or as a β -path defined as $\beta(t_1, t_2) = \gamma(t_2)$. Note also that any α -path can be considered as a quotient map of S^1 by the action of s_1 . Similarly, any β -path can be considered as a quotient map of S^1 by the action of s_2 .

With every α -path α , we can associate a *gluing map* $\pi_\alpha : \overline{\Delta}_\infty \rightarrow S^2$, a continuous map that restricts to an orientation-preserving homeomorphism between Δ_∞ and $S^2 - \alpha(S^1)$ and coincides with α on S^1 . The gluing map π_α thus defined is unique up to a homotopy relative to the unit circle. Similarly, we associate a gluing map $\pi_\beta : \overline{\Delta}_\infty \rightarrow S^2$ with every β -path β . Note that gluing maps are quotient maps of $\overline{\Delta}_\infty$ by the action of s_1 or s_2 on S^1 .

Let \mathcal{A} be a set of disjoint α -paths in the sphere. Being disjoint means that $\alpha(S^1) \cap \alpha'(S^1) = \emptyset$ for every pair of different α -paths $\alpha, \alpha' \in \mathcal{A}$. Define the set $\text{Im } \mathcal{A} \subset S^2$ as the union of $\alpha(S^1)$ for all $\alpha \in \mathcal{A}$. We define $\text{Im } \mathcal{B}$ for a set \mathcal{B} of disjoint β -paths in the same way. For every $\alpha \in \mathcal{A}$, fix a gluing map $\pi_\alpha : \overline{\Delta}_\infty \rightarrow S^2$. Define the *ungluing space* $\Upsilon_{\mathcal{A}}$ of \mathcal{A} as the subset of $\overline{\Delta}_\infty^{\mathcal{A}}$ given by the following condition. A point $\chi : \mathcal{A} \rightarrow \overline{\Delta}_\infty$ of $\overline{\Delta}_\infty^{\mathcal{A}}$ belongs to $\Upsilon_{\mathcal{A}}$ if the points $\pi_\alpha \circ \chi(\alpha) \in S^2$ are the same for all $\alpha \in \mathcal{A}$. In other terms, $\Upsilon_{\mathcal{A}}$ is the equalizer of the gluing maps $\pi_\alpha, \alpha \in \mathcal{A}$. Define the map $\pi_{\mathcal{A}} : \Upsilon_{\mathcal{A}} \rightarrow S^2$ as the map $\chi \mapsto \pi_\alpha \circ \chi(\alpha)$ (note that, by definition, $\pi_\alpha \circ \chi(\alpha)$ does not depend on α). Clearly, this map is continuous. Intuitively, $\Upsilon_{\mathcal{A}}$ is obtained from the sphere by ungluing all curves $\alpha(S^1), \alpha \in \mathcal{A}$, i.e. making topological circles




Fig. 1 A schematic picture of regluing

out of them. Note that a point $x \in S^2$ has only one preimage under the map $\pi_{\mathcal{A}}$ unless $x \in \alpha(S^1)$ for some $\alpha \in \mathcal{A}$. Similarly, we can define the ungluing space $\Upsilon_{\mathcal{B}}$ and the map $\pi_{\mathcal{B}} : \Upsilon_{\mathcal{B}} \rightarrow S^2$ for a set \mathcal{B} of disjoint β -paths.

Denote by $\Upsilon_{\mathcal{A}}^{\circ}$ the subset $\pi_{\mathcal{A}}^{-1}(S^2 - \text{Im } \mathcal{A})$ of the ungluing space $\Upsilon_{\mathcal{A}}$. Also, for every $\alpha \in \mathcal{A}$, we can define $S_{\alpha}^1 \subset \Upsilon_{\mathcal{A}}$ as the set of points $\chi \in \Upsilon_{\mathcal{A}}$ such that $\pi_{\mathcal{A}}(\chi) \in \alpha(S^1)$. There is a natural homeomorphism $h_{\alpha} : S^1 \rightarrow S_{\alpha}^1$ that takes a point $u \in S^1$ to the point $\chi \in S_{\alpha}^1$ defined as follows:

$$\chi(\alpha') = \begin{cases} u, & \alpha = \alpha', \\ \pi_{\alpha'}^{-1} \circ \alpha(u), & \alpha \neq \alpha'. \end{cases}$$

Thus the ungluing space splits into the union of the set $\Upsilon_{\mathcal{A}}^{\circ}$, which identifies canonically with $S^2 - \text{Im } \mathcal{A}$, and disjoint topological circles S_{α}^1 , which identify canonically with S^1 .

The Klein group V_4 acts on $\Upsilon_{\mathcal{A}}$. Namely, the action on $\Upsilon_{\mathcal{A}}^{\circ}$ is trivial, and the action on every circle S_{α}^1 identifies (through the homeomorphism h_{α}) with the standard action of V_4 on S^1 . It is not hard to prove that $\pi_{\mathcal{A}} : \Upsilon_{\mathcal{A}} \rightarrow S^2$ is a quotient map by the action of s_1 (warning: the action is not continuous; it still makes sense to talk about the quotient space by the orbit equivalence relation). Similarly, for a set \mathcal{B} of disjoint β -paths, the map $\pi_{\mathcal{B}} : \Upsilon_{\mathcal{B}} \rightarrow S^2$ is a quotient map by the action of s_2 .

We can now give a definition of a regluing. Here, we will only define a topological regluing of a set of disjoint simple curves, thus the word “regluing” will only be used in this sense until Sect. 5. Consider a set \mathcal{A} of disjoint α -paths and a set \mathcal{B} of disjoint β -paths in the sphere. A bijective continuous orientation-preserving map $\Phi : S^2 - \text{Im } \mathcal{A} \rightarrow S^2 - \text{Im } \mathcal{B}$ is called a regluing of \mathcal{A} into \mathcal{B} according to a given one-to-one correspondence between \mathcal{A} and \mathcal{B} if, for every $\alpha \in \mathcal{A}$ and the corresponding $\beta \in \mathcal{B}$, the map

$$\pi_{\beta}^{-1} \circ \Phi \circ \pi_{\alpha} : \pi_{\alpha}^{-1}(S^2 - \text{Im } \mathcal{A}) \rightarrow \pi_{\beta}^{-1}(S^2 - \text{Im } \mathcal{B})$$

extends to $S^1 \cup \pi_\alpha^{-1}(S^2 - \text{Im } \mathcal{A})$ so that the extension is continuous at all points of S^1 , and its restriction to S^1 is the identity.

Example 1 As an example, consider a branch of the function

$$j(z) = \sqrt{z^2 - 1}$$

on the sphere with the interval $[-1, 1]$ removed. Then j maps the set $\overline{\mathbb{C}} - [-1, 1]$ to the set $\overline{\mathbb{C}} - [-i, i]$ homeomorphically. Set $\alpha(t_1, t_2) = t_1$ and $\beta(t_1, t_2) = it_2$. The map j reglues α into β , for a suitable choice of the branch.

Proposition 1 Consider a regluing $\Phi : S^2 - \text{Im } \mathcal{A} \rightarrow S^2 - \text{Im } \mathcal{B}$. There exists a homeomorphism $\hat{\Phi} : \Upsilon_{\mathcal{A}} \rightarrow \Upsilon_{\mathcal{B}}$ that makes the following diagram commutative:

$$\begin{array}{ccccc} \Upsilon_{\mathcal{A}} & \xrightarrow{\pi_{\mathcal{A}}} & S^2 & \xleftarrow{\text{inclusion}} & S^2 - \text{Im } \mathcal{A} \\ \hat{\Phi} \downarrow & & & & \downarrow \Phi \\ \Upsilon_{\mathcal{B}} & \xrightarrow{\pi_{\mathcal{B}}} & S^2 & \xleftarrow{\text{inclusion}} & S^2 - \text{Im } \mathcal{B} \end{array}$$

Moreover, the homeomorphism $\hat{\Phi}$ commutes with the action of V_4 on the ungluing spaces.

Proof On $\Upsilon_{\mathcal{A}}^\circ$, we define $\hat{\Phi}$ as $\pi_{\mathcal{B}}^{-1} \circ \Phi \circ \pi_{\mathcal{A}}$. On S_α^1 , $\alpha \in \mathcal{A}$, define $\hat{\Phi}$ as $h_\beta \circ h_\alpha^{-1}$, where β is the β -path corresponding to the α -path α .

It is clear that $\hat{\Phi}$ is one-to-one. This map is continuous on $\Upsilon_{\mathcal{A}}^\circ$ by definition. The continuity on S_α^1 , $\alpha \in \mathcal{A}$, follows from the formula

$$\hat{\Phi} = \text{pr}_\beta^{-1} \circ \pi_\beta^{-1} \circ \Phi \circ \pi_\alpha \circ \text{pr}_\alpha$$

on $\Upsilon_{\mathcal{A}}^\circ$, where $\text{pr}_\alpha : \Upsilon_{\mathcal{A}} \rightarrow S^2$ is the projection $\chi \mapsto \chi(\alpha)$. Since the ungluing spaces are Hausdorff and compact, it follows that $\hat{\Phi}$ is a homeomorphism. Since $\hat{\Phi}$ acts as the identity under standard identifications of S_α^1 and S_β^1 with S^1 , the action of V_4 is preserved. □

For every α -path $\alpha : S^1 \rightarrow S^2$, define a β -path $\alpha^\#$ as follows: $\alpha^\#(t_1, t_2) = \alpha(t_2, t_1)$. Similarly, the formula $\beta^\#(t_1, t_2) = \beta(t_2, t_1)$ makes a β -path β into an α -path $\beta^\#$. For a set \mathcal{A} of α - or β -paths, we can form the set $\mathcal{A}^\# = \{\alpha^\# \mid \alpha \in \mathcal{A}\}$.

Proposition 2 Consider a regluing Φ of a set \mathcal{A} of disjoint α -paths into a set \mathcal{B} of disjoint β -paths.

- The map $\Phi : S^2 - \text{Im } \mathcal{A} \rightarrow S^2 - \text{Im } \mathcal{B}$ is a homeomorphism.
- The map Φ^{-1} is a regluing of $\mathcal{B}^\#$ into $\mathcal{A}^\#$.

Proof This follows immediately from the existence of the homeomorphism $\hat{\Phi}$ and the fact that it commutes with the action of V_4 on the ungluing spaces. \square

Let $f : S^2 \rightarrow S^2$ be a continuous map. Assume that a countable set \mathcal{A} of disjoint α -paths satisfies the following conditions:

- *Forward semi-invariance*: for any path $\alpha \in \mathcal{A}$, we have $f \circ \alpha \in \mathcal{A}$ or $f \circ \alpha(t_1, t_2) = f \circ \alpha(-t_1, t_2)$ for all $(t_1, t_2) \in S^1$. In the latter case, $f \circ \alpha(S^1)$ must be disjoint from $\text{Im } \mathcal{A}$. In the former case, the map $\pi_{\alpha \circ f}^{-1} \circ f \circ \pi_\alpha$ must be defined for all points of $\underline{\Delta}_\infty$ sufficiently close to S^1 , and must extend to a neighborhood of S^1 in $\underline{\Delta}_\infty$ so that the extension is continuous at all points of S^1 , and its restriction to S^1 is the identity.
- *Backward invariance*: we have $f^{-1}(\text{Im } \mathcal{A}) \subseteq \text{Im } \mathcal{A}$.

We say in this case that \mathcal{A} is *f-stable*.

Proposition 3 *If the set \mathcal{A} is f-stable, then there is a continuous map $\hat{f} : \Upsilon_{\mathcal{A}} \rightarrow \Upsilon_{\mathcal{A}}$ that makes the following diagram commutative:*

$$\begin{array}{ccc}
 \Upsilon_{\mathcal{A}} & \xrightarrow{\pi_{\mathcal{A}}} & S^2 \\
 \hat{f} \downarrow & & \downarrow f \\
 \Upsilon_{\mathcal{A}} & \xrightarrow{\pi_{\mathcal{A}}} & S^2
 \end{array}$$

Moreover, \hat{f} is equivariant with respect to the action of V_4 on $\Upsilon_{\mathcal{A}}$.

Proof Suppose that $f \circ \pi_{\mathcal{A}}(\chi) \notin \text{Im } \mathcal{A}$, then we set $\hat{f}(\chi) = \pi_{\mathcal{A}}^{-1} \circ f \circ \pi_{\mathcal{A}}(\chi)$. Now suppose that $f \circ \pi_{\mathcal{A}}(\chi) \in \alpha(S^1)$ for some $\alpha \in \mathcal{A}$. By the backward invariance, we necessarily have $\pi_{\mathcal{A}}(\chi) \in \alpha'(S^1)$ for some other $\alpha' \in \mathcal{A}$, hence $\chi(\alpha') \in S^1$. In this case, we can set $\hat{f}(\chi) = h_\alpha \circ \chi(\alpha')$. The continuity of \hat{f} at points of S^1_α follows from the forward semi-invariance and the formula

$$\hat{f} = \text{pr}_\alpha^{-1} \circ \pi_\alpha^{-1} \circ f \circ \pi_{\alpha'} \circ \text{pr}_{\alpha'}$$

on $\Upsilon_{\mathcal{A}}^\circ$. It is straightforward to check that \hat{f} commutes with the action of V_4 . \square

Our main construction is based on the following simple fact:

Theorem 2 *Suppose that $f : S^2 \rightarrow S^2$ is a continuous map, and \mathcal{A} is an f -stable set of disjoint α -paths. Let Φ be a regluing of \mathcal{A} into a set \mathcal{B} of disjoint β -paths. Then the map $g = \Phi \circ f \circ \Phi^{-1}$ extends to a continuous map from S^2 to S^2 . Moreover, the set \mathcal{B} is g -stable, and $\hat{g} = \hat{\Phi} \circ \hat{f} \circ \hat{\Phi}^{-1}$.*

Proof Consider the map $\hat{g} = \hat{\Phi} \circ \hat{f} \circ \hat{\Phi}^{-1}$, which takes $\Upsilon_{\mathcal{B}}$ to $\Upsilon_{\mathcal{B}}$. Since both $\hat{\Phi}$ and \hat{f} are equivariant under the action of V_4 , this map is also equivariant. It follows that it descends to a continuous map $g : S^2 \rightarrow S^2$. □

We would like to apply this theorem as follows. Let $f : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ be a rational function. For certain classes of rational functions f , there are natural ways to produce f -stable sets of paths. Then the corresponding map g is often a model for a new rational function. Note that the topological dynamics of g is very easy to understand in terms of the topological dynamics of f , because Φ is a topological conjugation except on $\text{Im}(\mathcal{A})$. A remarkable fact is that in many cases, the regluing Φ makes sense in a certain holomorphic category, so that the construction may actually produce a rational function g rather than just a continuous map.

Let X be a compact metric space, and \mathcal{S} a set of compact subsets of X . Recall that \mathcal{S} forms a null-sequence if for every $\varepsilon > 0$, there are only finitely many elements of \mathcal{S} , whose diameter exceeds ε . The following theorem is needed for the construction of topological models:

Theorem 3 *Let \mathcal{A} be a countable set of disjoint α -paths such that the sets $\alpha(S^1)$, $\alpha \in \mathcal{A}$, form a null-sequence. Then*

1. *the ungluing space $\Upsilon_{\mathcal{A}}$ embeds homeomorphically to S^2 ;*
2. *there exists a regluing of \mathcal{A} into some set \mathcal{B} of disjoint β -paths; moreover, one can arrange that $\beta(S^1)$, $\beta \in \mathcal{B}$ also form a null-sequence.*

The statement of the theorem may seem intuitively obvious (and it is in fact obvious for the case of finite \mathcal{A}). Note, however, that the set $\text{Im } \mathcal{A}$ may be everywhere dense in the sphere, and even have full measure. We will prove Theorem 3 in Sect. 6.3 using Moore’s theory. It is useful to know that the property of being a null-sequence is topological, and does not depend on a particular metric:

Proposition 4 *Let X be a compact metric space. A set \mathcal{S} of compact subsets of X forms a null-sequence if and only if for every open covering \mathcal{E} of X , there is a finite subset $\mathcal{S}' \subset \mathcal{S}$ such that every element of $\mathcal{S} - \mathcal{S}'$ is contained in an element of \mathcal{E} .*

In other terms, the number $\varepsilon > 0$ can be replaced with an open covering \mathcal{E} . The following proof is rather standard.

Proof We first prove the only if part. Let \mathcal{E} be an open covering of X , and ε its Lebesgue number. Recall that a Lebesgue number of \mathcal{E} is defined as a real number $\varepsilon > 0$ such that every set of diameter less than ε belongs to an element of \mathcal{E} . Set \mathcal{S}' to be the set of all elements of \mathcal{S} , whose diameter is at least ε . Then, by the Lebesgue number lemma, every element of $\mathcal{S} - \mathcal{S}'$ is contained in an element of \mathcal{E} .

Let us now prove the if part. Choose any $\varepsilon > 0$, and consider the covering \mathcal{E} of X by all $\varepsilon/2$ -balls. Then there is a finite subset $\mathcal{S}' \subset \mathcal{S}$ such that every set in $\mathcal{S} - \mathcal{S}'$ is contained in an element of \mathcal{E} . It follows that the diameter of any set in $\mathcal{S} - \mathcal{S}'$ does not exceed ε . □

Corollary 1 *Let X and Y be compact metric spaces, and $\phi : X \rightarrow Y$ a continuous map. If \mathcal{S} is a set of compact subsets in X that form a null-sequence, then the sets $\phi(A)$, $A \in \mathcal{S}$, form a null-sequence in Y .*

Proof Indeed, let \mathcal{E} be any open covering of Y . Consider the corresponding covering $\phi^*(\mathcal{E})$ of X . By Proposition 4, all elements of \mathcal{S} but finitely many are subsets of elements of $\phi^*(\mathcal{E})$. It follows that all $\phi(A)$, $A \in \mathcal{S}$, but finitely many are subsets of elements of \mathcal{E} . □

2.2 Regluing of branched coverings

The setting of branched coverings is most commonly used for topological discussions of rational functions. On one hand, branched coverings are objects of topological nature, and are much more flexible than holomorphic functions. On the other hand, they are nice objects and do not have pathologies of general continuous maps. This is why we want the regluing construction to fit into the contest of topological branched coverings.

Let $f : S^2 \rightarrow S^2$ be a branched covering. Consider a set \mathcal{A} of simple disjoint α -paths in the sphere such that $\alpha(S^1)$, $\alpha \in \mathcal{A}$, form a null-sequence. We say that \mathcal{A} is *strongly f -stable* if it is f -stable, and satisfies the following additional assumption: all critical points in $\text{Im } \mathcal{A}$ have the form $\alpha(0, 1)$, where $\alpha \in \mathcal{A}$ is a path such that $f \circ \alpha(t_1, t_2) = f \circ \alpha(-t_1, t_2)$ for all $(t_1, t_2) \in S^1$; moreover, these critical points are simple. Note that the null-sequence property is included into the notion of a strongly f -stable set of paths.

Theorem 4 *Let $f : S^2 \rightarrow S^2$ be a branched covering, and \mathcal{A} a strongly f -stable set of disjoint α -paths. Consider a regluing $\Phi : S^2 - \text{Im } \mathcal{A} \rightarrow S^2 - \text{Im } \mathcal{B}$ of \mathcal{A} into some set \mathcal{B} of disjoint β -paths. Then the map $g = \Phi \circ f \circ \Phi^{-1}$ extends to a branched self-covering of S^2 .*

We will prove this theorem in Sect. 6.4. In the statement of the theorem, the curves $\beta(S^1)$, $\beta \in \mathcal{B}$, will automatically form a null-sequence. Indeed, by

Theorem 3, there is a regluing Φ' of \mathcal{A} into some set \mathcal{B}' of disjoint β -paths such that $\beta'(S^1)$, $\beta' \in \mathcal{B}'$, form a null-sequence. Then $\Phi \circ (\Phi')^{-1}$ extends to the sphere as a continuous map. Moreover, it maps \mathcal{B}' to \mathcal{B} . By Corollary 1, the curves $\beta(S^1)$, $\beta \in \mathcal{B}$, must also form a null-sequence.

2.3 Topological regluing of quadratic polynomials

In this section, we will not say much new about the dynamics of quadratic polynomials. However, we can illustrate the idea of regluing using quadratic polynomials as an example.

Let \mathcal{R} be an external ray in the parameter plane of quadratic polynomials (we write quadratic polynomials in the form $p_c(z) = z^2 + c$, thus the parameter plane is the c -plane). Suppose that \mathcal{R} lands at a point c on the boundary of the Mandelbrot set. Suppose that the Julia set of $f = p_c$ is locally connected, and that all periodic points of f (except ∞) are repelling. The ray \mathcal{R} determines a pair of rays R_f^+ and R_f^- in the dynamical plane of f that land at the critical point 0 (for parameter values in the ray \mathcal{R} , these two rays crash into 0). Note that there may be more rays landing at 0, but the pair of rays R_f^+ , R_f^- is distinguished (i.e. determined by the choice of \mathcal{R}).

Fix any real number $\rho > 0$. Consider the α -path $\alpha_0 : S^1 \rightarrow \mathbb{C}$ in the dynamical plane of f defined as follows:

$$\alpha_0(t_1, t_2) = \begin{cases} R_f^+(t_1\rho), & t_1 > 0, \\ 0, & t_1 = 0, \\ R_f^-(-t_1\rho), & t_1 < 0. \end{cases}$$

Here the dynamical rays are parameterized by the values of the Green function, thus $R(t)$ stands for the point in the ray R , at which the Green function is equal to t . Then, for each $n \geq 0$, the multivalued function $f^{-n} \circ \alpha_0$ has 2^n branches, each being an α -path. All these paths are called *pullbacks* of α_0 under the iterates of f . Let \mathcal{A} denote the set of such pullbacks, including α_0 . Clearly, \mathcal{A} is strongly f -stable.

Now consider the quadratic polynomial $g = p_{c_0}$, where $c_0 = \mathcal{R}(2\rho)$ is the point on the external parameter ray \mathcal{R} with parameter 2ρ (the external parameter rays are parameterized by the value of the Green function at the critical value). This means that, in the dynamical plane of g , the value of the Green function at the critical value c_0 is equal to 2ρ . Therefore, the value of the Green function at the critical point 0 is equal to ρ . There are exactly two rays that are bounded and contain 0 in their closures (here by a ray we mean any gradient curve of the Green function). Denote these rays by R_g^+ and R_g^- . These two rays can also be parameterized by the values of the Green function, thus the parameter runs through the interval $(0, \rho)$. Let z^+ and z^- be the

landing points of the rays R_g^+ and R_g^- (these rays land because the angle of \mathcal{R} cannot be a rational number with an odd denominator, which is the only case where one of the rays R_g^+ and R_g^- can crash into a precritical point). Define the following β -path $\beta_0 : S^1 \rightarrow \mathbb{C}$ in the dynamical plane of g :

$$\beta_0(t_1, t_2) = \begin{cases} z^+, & t_2 = 1, \\ R_g^+(|t_1|\rho), & t_2 > 0, \\ 0, & t_2 = 0, \\ R_g^- (|t_1|\rho), & t_2 < 0, \\ z^-, & t_2 = -1. \end{cases}$$

Let \mathcal{B} denote the set of all pullbacks of β_0 , including β_0 .

There is a natural one-to-one correspondence between the sets of paths \mathcal{A} and \mathcal{B} . For any path $\alpha \in \mathcal{A}$, the point $\alpha(1, 0)$ belongs to a unique external ray of angle θ . There is a unique path $\beta \in \mathcal{B}$ such that the ray of angle θ crashes into $\beta(1, 0)$. We will make this path β correspond to the path α .

Theorem 5 *There exists a regluing Φ of \mathcal{A} into \mathcal{B} such that $g(x) = \Phi \circ f \circ \Phi^{-1}(x)$ at all points x , where the right-hand side is defined.*

Before we proceed with the proof of this theorem, we need to recall the definition of symbolic itineraries. The union Γ_f of the rays R_f^+ and R_f^- together with their common landing point 0 divides the complex plane into two connected components. For a point x not in $f^{-n}(\Gamma_f)$, we define $\sigma_f^n(x)$ to be 0 or 1 depending on whether or not $f^{on}(x)$ is separated from the critical value c of f by Γ_f . The sequence of numbers $\sigma_f^n(x)$, $n = 1, 2, \dots$ (which may be finite or infinite depending on whether or not x is eventually mapped to Γ_f) is called the *symbolic itinerary* of x . Similarly, we define Γ_g to be the union of $\{0\}$ and the external rays in the dynamical plane of g that crash into 0 (they have the same external angles as the rays R_f^+ and R_f^-). The definition of symbolic itineraries carries over to the dynamical plane of g , where we use Γ_g instead of Γ_f . In the dynamical plane of g , as well as in the dynamical plane of f , there cannot be two different points in the Julia set with the same symbolic itinerary. This is a basic Poincaré distance argument, see e.g. [9].

Proof of Theorem 5 Consider the complement U to the closure of $\text{Im } \mathcal{A}$. Since the closure of $\text{Im } \mathcal{A}$ contains the Julia set of f , the set U is an open subset of the Fatou set. Actually, U is the complement in the Fatou set to $\text{Im } \mathcal{A}$.

We first define the map Φ just on U . Let ϕ_f be the Böttcher parameterization for f , i.e. the inverse of the Böttcher coordinate function. Since the Fatou set of f contains no precritical points except for ∞ , the map ϕ_f is a holomorphic isomorphism between the unit disk Δ and the Fatou set of f

such that $\phi_f(z^2) = f \circ \phi_f(z)$ for all $z \in \Delta$. Similarly, let ϕ_g be the Böttcher parameterization for g . Note that ϕ_g is not everywhere defined on the unit disk, because the Fatou set of g contains the critical point 0 and its preimages. However, ϕ_g is well-defined on the complement to $\phi_f^{-1}(\text{Im } \mathcal{A})$ in the unit disk, and gives a holomorphic isomorphism between this complement and the complement to $\overline{\text{Im } \mathcal{B}}$ in the sphere. The set $\phi_f^{-1}(\text{Im } \mathcal{A})$ consists of countably many intervals of the form $[z, z/|z|]$. Set $\Phi = \phi_g \circ \phi_f^{-1}$. The map Φ is defined on U . Clearly, we have $g = \Phi \circ f \circ \Phi^{-1}$ on $\Phi(U)$. Note also that Φ preserves the values of the Green function.

It is easy to see that Φ extends continuously to each side of each curve $\alpha(S^1)$, $\alpha \in \mathcal{A}$. The extension preserves the values of the Green function. It follows that, for every $\alpha \in \mathcal{A}$ and the corresponding $\beta \in \mathcal{B}$, the function $\pi_\beta^{-1} \circ \Phi \circ \pi_\alpha$ extends to $S^1 \cup \pi_\alpha^{-1}(U)$ so that the extension is continuous at all points of S^1 , and its restriction to S^1 is the identity.

We now need to show that for any point $z \in \mathbb{C} - \text{Im } \mathcal{A}$ and any sequence $z_n \in U$ converging to z , the sequence $w_n = \Phi(z_n)$ converges to a well-defined point in the dynamical plane of g , and that this point does not depend on the choice of the sequence z_n . Since Φ is continuous on U , it suffices to assume that z belongs to the Julia set of f . Any limit point of the sequence w_n must have the same symbolic itinerary as z , therefore, this can only be one point. We denote this point by $\Phi(z)$, which is justified by the fact that Φ extends continuously to z . Note that, for different points z in the Julia set of f but not in $\text{Im } \mathcal{A}$, the points $\Phi(z)$ have different symbolic itineraries, and hence are different. This finishes the proof of the theorem. □

The reason for the proof shown above to be so simple is that we know a lot about topological dynamics of both f and g . However, we would like to use regluing to describe new kinds of topological dynamics, and that would be necessarily more complicated.

2.4 Topological models via regluing

First, we need to make the notion of topological model more precise. Define an (abstract) *topological model* as the collection of the following data:

- A branched covering $f : X \rightarrow X$, where X is a topological space homeomorphic to the 2-sphere.
- A compact fully invariant subset $J \subset X$, called the *Julia set* of f . The complement to the Julia set is called the *Fatou set*.
- A complex structure (i.e. a structure of a one-dimensional complex manifold, not necessarily connected) on the Fatou set such that f is holomorphic with respect to this structure.

The topological space X , the map f and the set J are called the *model space*, the *model map* and the *model Julia set*, respectively. Instead of referring to a topological model as (f, X, J) , we will sometimes simply say “topological model f ”. We will sometimes call X the *dynamical sphere* of f . Of course, any rational function is an abstract topological model. We say that two abstract topological models (f, X_f, J_f) and (g, X_g, J_g) are *equivalent* if there is a homeomorphism $\phi : X_f \rightarrow X_g$ conjugating f with g and such that $\phi(J_f) = J_g$ and $\phi|_{X_f - J_f}$ is holomorphic. We say that f *models* a rational function R if f is equivalent to R as an abstract topological model.

Although a rational function can be given by a very explicit formula, it may be very hard to see its dynamics from this formula. Even drawing an accurate picture requires a lot of computational power. However, we need a *dynamically explicit* description of rational functions. In other terms, we want to find explicit topological models for rational functions. The notion of abstract topological model formalizes this goal, except for the notion of being (dynamically) explicit, which is, of course, an informal notion. The situation is similar to finding explicit solutions of algebraic equations—the notion of solution is rigorously defined, but the notion of being explicit is informal, and depends on the taste, philosophy, etc.

There are several important combinatorial constructions that modify or combine topological models into new topological models. Among the most well-known are matings and captures. Let us now define another combinatorial operation on topological models that uses regluing. There are interesting relationships between matings, captures and regluings, which we may discuss elsewhere.

Consider a topological model (f, X, J) with a hyperbolic Fatou set (i.e. the complex 1-manifold $X - J$ is of hyperbolic type), and a strongly f -stable set \mathcal{A} of disjoint α -paths in X . We will now assume that \mathcal{A} is *non-wandering*: for every $\alpha \in \mathcal{A}$, there exists $n > 0$ such that $f^{on} \circ \alpha \notin \text{Im } \mathcal{A}$. In particular, $\alpha(0, 1)$ is eventually mapped to a critical point of f . Under this assumption, we will define another topological model using regluing.

Define an *accumulation point* of \mathcal{A} as a point $x \in X$ such that every open neighborhood of x intersects infinitely many curves $\alpha(S^1)$, $\alpha \in \mathcal{A}$. All accumulation points of \mathcal{A} belong to the Julia set of f . Indeed, since $f : X - J \rightarrow X - J$ is a holomorphic self-map of a hyperbolic complex 1-manifold, backward f -orbits of critical points cannot accumulate in $X - J$ (see e.g. [9]). Since the curves $\alpha(S^1)$, $\alpha \in \mathcal{A}$, form a null-sequence, all accumulation points of \mathcal{A} are also accumulation points of the backward orbits of critical points.

There exists a regluing $\Phi : X - \text{Im } \mathcal{A} \rightarrow Y - \text{Im } \mathcal{B}$ of \mathcal{A} into some set \mathcal{B} of disjoint β -paths in a topological sphere Y . Moreover, the curves $\beta(S^1)$, $\beta \in \mathcal{B}$, form a null-sequence. We set g to be the continuous extension of $\Phi \circ f \circ \Phi^{-1}$, which exists by Theorem 2. Since f is a branched covering, and

\mathcal{A} is strongly f -stable, the map g is also a branched covering by Theorem 4. Define the Fatou set of g as the set of all points $y \in Y$ such that, for some nonnegative integer n , we have $g^{on}(y) \notin \text{Im } \mathcal{B}$ (thus Φ^{-1} is defined at this point) and $\Phi^{-1}(g^{on}(y)) \in X - J$. Clearly, if this condition is satisfied for one particular n , then it also holds for all bigger n . Therefore, the Fatou set of g thus defined is fully invariant.

It remains to define a complex structure on the Fatou set of g invariant under g . Take a point y in the Fatou set of g and its iterated image $y' = g^{on}(y)$ such that $y' \notin \text{Im } \mathcal{B}$ and $x' = \Phi^{-1}(y')$ is in the Fatou set of f . Since \mathcal{A} does not accumulate in the Fatou set of f , there is a neighborhood U of x' disjoint from $\text{Im } \mathcal{A}$, and a holomorphic embedding $\xi : U \rightarrow \mathbb{C}$ such that $\xi(x') = 0$. Note that $\Phi : U \rightarrow \Phi(U)$ is a homeomorphism. Therefore, $\xi \circ \Phi^{-1}$ is an embedding of the neighborhood $\Phi(U)$ of y' into \mathbb{C} . Finally, if k is the local degree of g^{on} at y , then we can define a local complex coordinate near y as a branch of $\sqrt[k]{\xi \circ \Phi^{-1} \circ g^{on}}$. We have now defined a complex coordinate near every point of the Fatou set of g . It is easy to check that all transition functions are holomorphic, and that g is holomorphic with respect to the obtained complex structure on the Fatou set.

We have defined a topological model g . This topological model will be called the model obtained from (f, X, J) by regluing of \mathcal{A} . It is not hard to check that J lifts to the ungluing space $\Upsilon_{\mathcal{A}}$ as a compact subset \hat{J} fully invariant under both \hat{f} and the action of V_4 . Moreover, the set $\hat{\Phi}(\hat{J})$ is precisely the lift of the Julia set of g to the ungluing space $\Upsilon_{\mathcal{B}}$.

3 Boundary points of type C hyperbolic components

In this section, we discuss some general properties of quadratic rational functions on the boundaries of type C hyperbolic components, preparing for the proof of Theorem 1.

3.1 Equicontinuous families and holomorphic motion

Below, we recall a standard fact about equicontinuous families widely used in complex dynamics:

Proposition 5 *Let Λ be a topological space, and X a metric space. Consider an equicontinuous family \mathcal{F} of maps from Λ to X . Let $S \subset \Lambda$ be a subset, and $v : \Lambda \rightarrow X$ a continuous map such that $v(\lambda) \in \mathcal{F}(\lambda)$ for all $\lambda \in S$. Then for all $\lambda^* \in \overline{S}$, we have*

$$v(\lambda^*) \in \overline{\mathcal{F}(\lambda^*)}.$$

Proof Assume the contrary: $d(v(\lambda^*), \mathcal{F}(\lambda^*)) = \varepsilon > 0$, where d denotes the distance in X . There is a neighborhood V' of λ^* in Λ such that

$$d(v(\lambda), v(\lambda^*)) < \frac{\varepsilon}{2}$$

for all $\lambda \in V'$. This follows from the continuity of v . On the other hand, there is a neighborhood V'' of λ^* such that

$$d(f(\lambda), f(\lambda^*)) < \frac{\varepsilon}{2}$$

for all $\lambda \in V''$ and all $f \in \mathcal{F}$. This follows from the equicontinuity of \mathcal{F} . Therefore, for every $\lambda \in V' \cap V''$ and every $f \in \mathcal{F}$, we have

$$\begin{aligned} d(v(\lambda^*), f(\lambda^*)) &\leq d(v(\lambda^*), v(\lambda)) + d(v(\lambda), f(\lambda)) + d(f(\lambda), f(\lambda^*)) \\ &< d(v(\lambda), f(\lambda)) + \varepsilon. \end{aligned}$$

Take $\lambda \in V' \cap V'' \cap S$ and $f \in \mathcal{F}$ such that $v(\lambda) = f(\lambda)$. Then $d(v(\lambda^*), f(\lambda^*)) < \varepsilon$, a contradiction. \square

Let Λ be a complex analytic manifold, and A a set. Recall that a *holomorphic motion* over Λ is a map $\mu : \Lambda \times A \rightarrow \overline{\mathbb{C}}$ such that $\mu(\lambda, a) \neq \mu(\lambda, b)$ for $a \neq b$ and the map $\mu_a : \lambda \mapsto \mu(\lambda, a)$ is holomorphic for every $a \in A$. We do not require that $A \subset \overline{\mathbb{C}}$ and that $a \mapsto \mu(\lambda_0, a)$ is the identity for some $\lambda_0 \in \Lambda$, although these conditions are usually included into a definition. Thus we use the term “holomorphic motion” in a slightly more general sense. The following well-known fact is very simple but important (see e.g. [7]):

Theorem 6 *Let Λ be a Riemann surface and $\mu : \Lambda \times A \rightarrow \overline{\mathbb{C}}$ a holomorphic motion. Then the family of functions μ_a , $a \in A$, is equicontinuous.*

Proof If A is finite, then the statement is obvious. Suppose that A is infinite, and take three different points $a_1, a_2, a_3 \in A$. We can use the following generalization of Montel’s theorem: if a family of holomorphic functions on Λ is such that the graphs of all functions in the family avoid the graphs of three different holomorphic functions, and these three graphs are disjoint, then the family is equicontinuous. In our case, we can take μ_{a_i} , $i = 1, 2, 3$. These three holomorphic functions have disjoint graphs, and the graph of any function μ_a , $a \neq a_1, a_2, a_3$, is disjoint from the graphs of μ_{a_i} . Thus the family of functions μ_a is equicontinuous. \square

The following well-known theorem is proved in [7]:

Theorem 7 *Let Λ be a Riemann surface, and $\mu : \Lambda \times A \rightarrow \overline{\mathbb{C}}$ a holomorphic motion. Suppose that $A \subset \overline{\mathbb{C}}$ and that $\mu(\lambda_0, a) = a$ for some $\lambda_0 \in \Lambda$ and all $a \in A$. Then μ extends to a holomorphic motion $\overline{\mu} : \Lambda \times \overline{A} \rightarrow \overline{\mathbb{C}}$, and, for every $\lambda \in \Lambda$, the map $a \mapsto \overline{\mu}(\lambda, a)$ from \overline{A} to $\overline{\mathbb{C}}$ is quasi-symmetric.*

Using this theorem, we can prove the following (cf. e.g. [1]):

Proposition 6 *Consider a holomorphic motion μ satisfying the assumptions of Theorem 7. Assume that A is an open subset of $\overline{\mathbb{C}}$. Consider a continuous function $v : \Lambda \rightarrow \overline{\mathbb{C}}$ and the subset O of Λ consisting of all $\lambda \in \Lambda$ such that $v(\lambda) \in \mu(\lambda, A)$. Then O is open. Moreover, $v(\lambda^*) \in \partial\mu(\lambda, A)$ if $\lambda^* \in \partial O$.*

Proof Consider the point λ_0 from Theorem 7. We can assume without loss of generality that $\lambda_0 \in O$. Then $v(\lambda_0)$ is some point $a_0 \in A$. Suppose that $a_0 \neq \infty$. Let $\alpha : S^1 \rightarrow A$ be a small loop around a_0 . Then we have

$$I(\lambda_0) = \int_{S^1} \frac{d\alpha(t)}{\alpha(t) - a_0} = 2\pi i.$$

Set ε to be the minimal spherical distance between a_0 and $\alpha(t)$. There is an open neighborhood V of λ_0 such that the distance between $v(\lambda)$ and $\mu(\lambda, \alpha(t))$ is bigger than $\varepsilon/2$ for all $\lambda \in V$ and all $t \in S^1$. This follows from the continuity of v and equicontinuity of $\mu(\cdot, a)$, $a \in A$. We can also assume that $\mu(\lambda, \alpha(t)) \neq \infty$ for all $\lambda \in V$ and $t \in S^1$. Then the integral

$$I(\lambda) = \int \frac{d\mu(\lambda, \alpha(t))}{\mu(\lambda, \alpha(t)) - v(\lambda)}$$

(with respect to t) is a well-defined and continuous function on V . Since the possible values of this integral are discrete, we must have $I(\lambda) = 2\pi i$ for all $\lambda \in V$. Therefore, $v(\lambda) \in \mu(\lambda, A)$ for such λ , and $V \subset O$. Thus we have proved that O is open.

Suppose now that $\lambda^* \in \partial O$. Then $v(\lambda^*) \in \overline{\mu(\lambda^*, A)}$ by Proposition 5. On the other hand, we have $v(\lambda^*) \notin \mu(\lambda^*, A)$ because $\lambda^* \notin O$. Therefore, $v(\lambda^*) \in \partial\mu(\lambda, A)$. □

3.2 Parameter curves

Quadratic rational functions that are conjugate by a Möbius transformation have the same dynamical properties. Therefore, one wants to parameterize conjugacy classes, choosing one (or finitely many) particular representative(s) from every conjugacy class. There are many different ways to do this parameterization, see e.g. [8, 15]. For our purposes, it will be convenient to do the following: send the two critical points of a rational function f to 0 and ∞ by

a suitable Möbius transformation. If ∞ is fixed, then we can reduce f to the form $p_c : z \mapsto z^2 + c$. If ∞ is not fixed, then we can send a preimage of ∞ to 1 (unless 0 is a critical point mapping to ∞ , i.e. f is conjugate to $z \mapsto 1/z^2$), thus f will have the form

$$R_{a,b}(z) = \frac{az^2 - b}{z^2 - 1}.$$

In any case, f is Möbius conjugate to p_c , or to $R_{a,b}$, or to $1/z^2$.

We will now consider the following algebraic curves in \mathbb{C}^2 :

$$\mathcal{V}_k = \{(a, b) \mid R_{a,b}^{\circ k-1}(\infty) = 1\}, \quad k = 2, 3, \dots$$

These are complex one-dimensional slices of the parameter space of quadratic rational functions. These slices correspond to simple (periodic) types of behavior of one critical point (note that $R_{a,b}(1) = \infty$ so that for all $(a, b) \in \mathcal{V}_k$, the critical point ∞ of the function $R_{a,b}$ is periodic of period k).

We will identify pairs $(a, b) \in \mathcal{V}_k$ with the corresponding rational functions $R_{a,b}$. For every $(a, b) \in \mathcal{V}_k$, let $\Omega_{a,b}$ denote the immediate basin of the super-attracting fixed point ∞ of the rational function $R_{a,b}^{\circ k}$. We will sometimes write Ω_F instead of $\Omega_{a,b}$ if $F = R_{a,b}$. Define the set

$$\mathcal{B}_k = \{(a, b) \in \mathcal{V}_k \mid 0 \in R_{a,b}^{om}(\Omega_{a,b}), m \geq 0\}.$$

This set consists of all parameter values such that the critical point 0 is in the immediate basin of the cycle of ∞ . Thus \mathcal{B}_k is the union of all type B hyperbolic components in \mathcal{V}_k . Define the set

$$\Lambda_k = \mathcal{V}_k - \overline{\mathcal{B}_k}.$$

This is a one-dimensional complex manifold (for smoothness, see e.g. [14, 16]).

Recall that a function $R_{a,b} \in \Lambda_k$ is hyperbolic of type C if $R_{a,b}^{om}(0) \in \Omega_{a,b}$ for some $m > 0$. The set of type C hyperbolic functions is open by Proposition 6.

3.3 Notation needed for the proof of Theorem 1

We will use the following notation throughout the proof of Theorem 1: Let $H \subset \Lambda_k$ be a hyperbolic component of type C, and $f \in \partial H$. Note that the boundary is taken in Λ_k , so that the boundaries of type B components are automatically excluded. Set $\Omega = \Omega_f$. Also, let h be the center of the hyperbolic component H , i.e. the unique critically finite map in H . There is a positive integer k' such that for any parameter value $(a, b) \in H$, we have $R_{a,b}^{\circ k'}(0) \in \Omega_{a,b}$, and k' is the minimal integer with this property.

Let Δ denote the unit disk $\{|z| < 1\}$. There is a holomorphic motion

$$\mu : \Lambda_k \times \Delta \rightarrow \overline{\mathbb{C}}$$

such that $\mu((a, b), z)$ is the point in $\Omega_{a,b}$, whose Böttcher coordinate is equal to z . By Theorem 7, this holomorphic motion extends to a holomorphic motion

$$\overline{\mu} : \Lambda_k \times \overline{\Delta} \rightarrow \overline{\mathbb{C}}.$$

By Proposition 6, we have $f^{ok'}(0) \in \partial\overline{\mu}(\lambda_0, \Delta) = \partial\Omega$, where $\lambda_0 \in \Lambda_k$ is the parameter value corresponding to f . Therefore, $f(0)$ is on the boundary of some Fatou component V containing a point v such that $f^{ok'-1}(v) = \infty$.

Note that there may be several Fatou components containing $f(0)$ on their boundary and eventually mapping to Ω . The choice of the Fatou component V is determined by the choice of the hyperbolic component H : as the parameter value enters H , the critical value corresponding to $f(0)$ enters a Fatou component corresponding to V (here “corresponding” means “included into the holomorphic motion of”).

3.4 Accessibility and non-recurrence

The following proposition shows that the Fatou components of f do not have topological pathologies (cf. [1]):

Proposition 7 *The boundary of Ω is locally connected. In particular, every boundary point is accessible from Ω .*

Proof Let λ_0 be the parameter value corresponding to f . Then the function $\overline{\mu}_f : z \mapsto \overline{\mu}(\lambda_0, z)$ is a quasi-symmetric homeomorphism between $\overline{\Delta}$ and $\overline{\Omega}$, by Theorem 7. It follows that the boundary of Ω is locally connected. \square

Since V is a pullback of Ω , the boundary of V is also locally connected. In particular, $f(0) \in \partial V$ is accessible from V . The following proposition shows that the dynamical properties of f are rather simple:

Proposition 8 *The critical point 0 of f is non-recurrent.*

Proof Note that all limit points of the orbit of 0 belong to the forward orbit of $\partial\Omega$ under f . Therefore, if 0 is recurrent, then $0 \in f^{ol}(\partial\Omega)$ for some $l = 0, \dots, k - 1$. In other terms, $0 = f^{ol}(\overline{\mu}(\lambda_0, z))$, where z is a point on the unit circle, and λ_0 is the parameter value corresponding to f .

Consider the holomorphic function $\nu(\lambda) = R_\lambda^{ol}(\overline{\mu}(\lambda, z))$ on Λ_k . (Recall that R_λ is the rational function $R_{a,b}$ corresponding to the point $\lambda = (a, b) \in \Lambda_k$.) This function vanishes at the point λ_0 . However, ν is not identically

equal to zero, because e.g. for $h \in H$, the critical point is not in the orbit of $\partial\Omega_h$. Therefore, we can choose a small loop $\gamma : S^1 \rightarrow \Lambda_k$ around λ_0 such that $\nu \circ \gamma$ loops around 0 (i.e. the algebraic number of full turns is nonzero). Take $\tilde{z} \in \Delta$ sufficiently close to z . Then the function $\tilde{\nu}(\lambda) = R_{\lambda}^{ol}(\overline{\mu}(\lambda, \tilde{z}))$ is uniformly close to ν . In particular, $\tilde{\nu} \circ \gamma$ loops around 0. Therefore, there exists a parameter value λ_1 inside γ such that $\tilde{\nu}(\lambda_1) = 0$. This means that 0 lies in $R_{\lambda_1}^{ol}(\Omega_{\lambda_1})$, i.e. $\lambda_1 \in \mathcal{B}_k$, a contradiction. \square

3.5 Restatement of Theorem 1

In this section, we will restate Theorem 1, and give some details on the particular set of paths that was mentioned but not defined in the statement of Theorem 1. We also need to introduce some more notation. Consider a simple path $\alpha_{-1} : [0, 1] \rightarrow \overline{V}$ such that $\alpha_{-1}(0) = f(0)$, $\alpha_{-1}(1) = \nu$, and $\alpha_{-1}(0, 1] \subset V$. The existence of such path follows from Proposition 7. There is an α -path $\alpha_0 : S^1 \rightarrow \overline{\mathbb{C}}$ such that $f \circ \alpha_0(t_1, t_2) = \alpha_{-1}(|t_1|)$. This α -path is unique up to the change of variables $t_1 \rightarrow -t_1$. Let \mathcal{A} be the set of all pullbacks of α_0 under iterates of f , including α_0 .

Proposition 9 *The curves $\alpha(S^1)$, $\alpha \in \mathcal{A}$ form a null-sequence.*

Proof Since the critical point 0 is non-recurrent, there exists a neighborhood U of $\alpha_0(S^1)$ that does not intersect the post-critical set. Choose a smaller neighborhood U' of $\alpha_0(S^1)$ that is compactly contained in U . By the Koebe distortion theorem, the set of pullbacks of U' forms a null-sequence. It follows that the curves $\alpha(S^1)$, $\alpha \in \mathcal{A}$, also form a null-sequence. \square

We can conclude that there exists a regluing Φ of \mathcal{A} into some set of disjoint β -paths. Moreover, the function $g = \Phi \circ f \circ \Phi^{-1}$ is well-defined as a topological model, since \mathcal{A} is strongly f -stable and non-wandering (both properties follow immediately from the definition of \mathcal{A}). We can now give a precise statement, from which Theorem 1 follows:

Theorem 8 *The map g is equivalent as a topological model to the critically finite hyperbolic rational function $h \in H$.*

Theorem 8 implies Theorem 1. Indeed, if $g = \Phi \circ f \circ \Phi^{-1}$ is conjugate to h , then $h = \Psi \circ f \circ \Psi^{-1}$ for some topological regluing Ψ . It follows that $f = \Psi^{-1} \circ h \circ \Psi$. Note that Ψ^{-1} is also a regluing.

Thus it remains to prove Theorem 8. We first prove that g is Thurston equivalent to h . By a theorem of Mary Rees [13], Thurston equivalence to a hyperbolic rational function implies semi-conjugacy. We will recall the proof of this theorem. What remains is to prove that all fibers are trivial. Several

ideas for this part were also taken from [13]. Overall, the argument is rather simple, but we need to know from the very beginning that g is well-defined as a branched covering. Here we use results of Sect. 6 on the existence of topological regluing.

4 The proof of Theorem 1

In this section, we prove Theorem 1. We use notation introduced in Sects. 3.5 and 3.3.

4.1 Backward stability

Recall the following theorem of Mañe [4, 6]:

Theorem 9 (Backward stability) *Let F be a rational function with the Julia set J . Suppose that there are no recurrent critical points in J . Then, for every $\varepsilon, \varepsilon' > 0$, there exist a positive real number $\delta(\varepsilon)$ (depending only on ε), and a positive integer $n_0(\varepsilon, \varepsilon')$ (depending on ε and ε') such that for $x, y \in J$*

1. *if $d(x, y) < \delta(\varepsilon)$, then for every $n \geq 0$ and for every $x' \in F^{-n}(x)$, there is a point $y' \in F^{-n}(y)$ such that $d(x', y') < \varepsilon$,*
2. *if $d(x, y) < \delta(\varepsilon)$ and $n \geq n_0(\varepsilon, \varepsilon')$, then for every $x' \in F^{-n}(x)$, there is a point $y' \in F^{-n}(y)$ such that $d(x', y') < \varepsilon'$.*

Here d is any metric on $\overline{\mathbb{C}}$ compatible with the topology.

By Proposition 8, this theorem is applicable to the function f . From the backward stability of f on its Julia set J_f , we can deduce the backward stability of g on its Julia set J_g :

Proposition 10 (Backward stability of g on J_g) *The map g is backwards stable on J_g , in the sense of Theorem 9.*

Proof Let \hat{f} be the lift of f to the ungluing space Υ_A , and \hat{J}_f the lift of the Julia set J_f . It is easy to see that \hat{f} is backwards stable on \hat{J}_f . This follows from the backward stability of f on J_f and the fact that the fibers of the projection $\pi_A : \hat{J}_f \rightarrow J_f$ form a null-sequence. Since backward stability is a topological property, \hat{g} is backwards stable on \hat{J}_g . It follows that g is backwards stable on J_g . □

4.2 Thurston equivalence

The map g is critically finite. Indeed, the critical values of g are $\Phi \circ f(\infty)$ and $\Phi(v)$. The forward orbit of $\Phi \circ f(\infty)$ under g is the Φ -image of the

forward orbit of $f(\infty)$ under f . Similarly, the forward orbit of $\Phi(v)$ under g is the Φ -image of the forward orbit of v under f . It remains to note that both $f(\infty)$ and v have finite forward orbits under f . In this section, we will prove that the map g is Thurston equivalent to h .

We first recall the notion of Thurston equivalence. A branched self-covering of the sphere with finite post-critical set is sometimes called a *Thurston map*. Recall that the post-critical set is the union of forward orbits of all critical values. Two Thurston maps F and G with post-critical sets P_F and P_G , respectively, are called *Thurston equivalent* if there exist homeomorphisms $\phi, \psi : S^2 \rightarrow S^2$ that make the following diagram commutative

$$\begin{array}{ccc}
 (S^2, P_F) & \xrightarrow{F} & (S^2, P_F) \\
 \psi \downarrow & & \downarrow \phi \\
 (S^2, P_G) & \xrightarrow{G} & (S^2, P_G)
 \end{array}$$

and such that ϕ and ψ are homotopic relative to P_F through homeomorphisms, in particular, $\phi|_{P_F} = \psi|_{P_F}$. The following are well-known useful criteria of Thurston equivalence:

Theorem 10 *Suppose that $F_t, t \in [0, 1]$, is a continuous family of Thurston maps of degree 2 such that the number of points in P_{F_t} does not change with t . Then F_0 is Thurston equivalent to F_1 .*

Let C_F denote the set of critical points of f .

Theorem 11 *Let Z be a compact connected locally connected subset of S^2 such that $S^2 - Z$ is connected and dense. Suppose that quadratic Thurston maps F and G are such that $F = G$ on Z and $P_F \cup C_F = P_G \cup C_G \subset Z$. Then F and G are Thurston equivalent.*

For completeness, we sketch the proofs of these theorems.

Proof of Theorem 10 It suffices to prove that F_t is Thurston equivalent to $F_{t'}$ provided that t is close to t' . The branched coverings F_t and $F_{t'}$ are then uniformly close, therefore, their post-critical sets are also close (this follows from the fact that the (weighted) number of critical points in a disk can be computed as a winding number). In particular, these maps are conjugate on their post-critical sets. Moreover, we can choose a homeomorphism $\phi : S^2 \rightarrow S^2$ that is close to the identity, maps P_{F_t} to $P_{F_{t'}}$ and conjugates the dynamics of F_t on P_{F_t} with the dynamics of $F_{t'}$ on $P_{F_{t'}}$. The multivalued function $F_{t'}^{-1} \circ \phi \circ F_t$ has two single valued branches. Indeed, every critical value of

$\phi \circ F_t$ is a critical value of $F_{t'}$, and the only $F_{t'}$ -preimage of this critical value is the corresponding critical point, since the map is quadratic. Thus, whenever $\phi(F_t(z))$ is a ramification point of $F_{t'}^{-1}$, the function $F_{t'}^{-1} \circ \phi \circ F_t$ can be written as $\sqrt{u^2}$ in some local coordinate. It follows that this function has no ramification points, hence it splits into two single valued branches. One of the branches is close to the identity; denote this branch by ψ . The homeomorphisms ϕ and ψ thus constructed provide a Thurston equivalence between F_t and $F_{t'}$ (they are homotopic relative to the post-critical set because they are both close to the identity). \square

Proof of Theorem 11 We can assume without loss of generality that $F = G$ on some neighborhoods of all critical points. This can be arranged e.g. by a small variation of G near its critical points, without changing the post-critical set.

By the same argument as in the proof of Theorem 10, the multivalued function $G^{-1} \circ F$ splits into two single valued branches. Near every critical point, one of the branches of $G^{-1} \circ F$ is the identity. It follows that there is a branch ϕ of $G^{-1} \circ F$ that restricts to the identity on Z . Indeed, the branches cannot switch outside of the critical points, and $z \in G^{-1}(F(z))$ for all $z \in Z$. There exists a continuous map $\gamma : \overline{\Delta} \rightarrow S^2$ that restricts to a holomorphic homeomorphism between Δ and $S^2 - Z$ (the set Z is infinite because any quadratic map has two different critical values). It is not hard to see (with the help of Carathéodory's theory) that $\gamma^{-1} \circ \phi \circ \gamma$ extends continuously to a self-homeomorphism of $\overline{\Delta}$ that is the identity on the unit circle. Such homeomorphism is isotopic to the identity relative to the unit circle (i.e. through homeomorphisms that are the identity on S^1). It follows that ϕ is isotopic to the identity through homeomorphisms of S^2 , whose restrictions to Z , in particular, to the post-critical set $P_F = P_G$, are the identity. The theorem follows. \square

Consider a simple curve in the parameter space Λ_k that connects f to h and lies entirely in H except for the endpoint f . Let $f_t, t \in [0, 1]$ be the corresponding one-parameter family of rational functions so that $f_0 = f$ and $f_1 = h$. Consider a point v_t depending continuously on $t \in [0, 1]$ and satisfying the following properties:

$$v_0 = v, \quad f_t^{\circ k'-1}(v_t) = \infty.$$

Then $v_1 = h(0)$. Define a quadratic rational function Q_t by the property that the critical values of Q_t coincide with $f_t(\infty)$ and v_t . We can even assume that the corresponding critical points of Q_t are ∞ and 0 , respectively (explicit formulas for such functions are very easy to obtain; they are given in Sect. 5). There exists a continuous family of α -paths $\alpha_t, t \in [0, 1)$

connecting the two preimages of v_t under f_t and having the property that $\alpha_t(-t_1, t_2) = -\alpha_t(t_1, t_2)$. Additionally, we can assume that α_t converge to the constant path 0 uniformly as $t \rightarrow 1$. Moreover, this continuous family can be chosen in such a way that for $t = 0$, we obtain the same α -path α_0 as that introduced in Sect. 3.5. (Recall that α_0 and all its pullbacks form the set \mathcal{A} of α -paths that is used in our regluing construction.) The multivalued analytic function $Q_t^{-1} \circ f_t$ has a holomorphic branch on the complement to $\alpha_t(S^1)$. Denote this branch by j_t . The choice of particular branch is not important; however, we need that j_t depend continuously on t . This can obviously be arranged.

Set $q_t = j_t \circ Q_t$. This function is defined on the complement to $Q_t^{-1}(\alpha_t(S^1))$, i.e. on the complement to a pair of disjoint simple curves. Intuitively, the function q_t is obtained from f_t by regluing the path α_t , or, to be more precise, we have $q_t = j_t \circ f_t \circ j_t^{-1}$ wherever the right-hand side is defined (however, q_t is defined on a larger set). Note that q_t is always defined and analytic in a neighborhood of 0, because $Q_t(0) = v_t$ avoids the curve $\alpha_t(S^1)$. In particular, 0 is a critical point of q_t . Another critical point is $j_t(\infty) = \infty$. Moreover, the critical orbits of q_t are finite and of constant size. More precisely, we have

$$q_t^{\circ k'}(0) = j_t \circ f_t^{\circ k'-1}(v_t) = \infty, \quad q_t^{\circ k}(\infty) = j_t \circ f_t^{\circ k}(\infty) = \infty.$$

In other words, the orbits of 0 and ∞ under q_t have the same dynamics as the orbits of the same points under h . The relations given above can be easily proved using that $q_t = j_t \circ Q_t$ and that $Q_t \circ j_t = f_t$ wherever the left-hand side is defined. The construction of the map q_t will reappear in Sect. 5, where more details can be found.

The maps q_t are critically finite but they are not Thurston maps because of discontinuities. However, one can “approximate” q_t by branched coverings \hat{q}_t that differ from q_t only in a small neighborhood of $Q_t^{-1}(\alpha_t(S^1))$. Of course, in a small neighborhood of $Q_t^{-1}(\alpha_t(S^1))$, we cannot hope to make the maps \hat{q}_t uniformly close to q_t , thus the word “approximate” comes in the quotation marks. However, we can do the following. Choose a sufficiently small Jordan neighborhood D of $\alpha_t(S^1)$. Then $Q_t^{-1}(D)$ is a pair of disjoint Jordan disks D_1 and D_2 . The boundaries $\partial D_1, \partial D_2$ get mapped to the same Jordan curve by q_t , moreover, q_t acts as an orientation-preserving homeomorphism on $\partial D_i, i = 1, 2$. Therefore, it is possible to extend q_t homeomorphically inside $D_1 \cup D_2$. The extension thus obtained is our map \hat{q}_t . We can choose \hat{q}_t to vary continuously with t . Additionally, we can arrange that $\hat{q}_1 = h$; in any case, \hat{q}_1 is Thurston equivalent to h . Thus \hat{q}_t is a continuous family of Thurston maps, whose post-critical sets are of constant cardinality. By Theorem 10, we obtain

Lemma 1 *The map \hat{q}_0 is Thurston equivalent to $\hat{q}_1 = h$.*

What remains to prove is the following

Lemma 2 *The maps \hat{q}_0 and g are Thurston equivalent.*

Proof Recall that $g = \Phi \circ f \circ \Phi^{-1}$ wherever the right-hand side is defined. Here Φ is a regluing of the set \mathcal{A} of α -paths consisting of the path α_0 and all its pullbacks under $f_0 = f$. The corresponding set of β -paths in the dynamical sphere of g is \mathcal{B} . Let β_0 be the β -path corresponding to the α -path α_0 .

Consider the map $j_0 \circ \Phi^{-1}$. It is defined on the complement to $\text{Im } \mathcal{B}$, it establishes a homeomorphism between $S^2 - \text{Im } \mathcal{B}$ and $j_0(S^2 - \text{Im } \mathcal{A})$, and extends continuously to $\beta_0(S^1)$. Denote the extension by Ψ^{-1} . Then Ψ is a homeomorphism between the complement to $j_0(\text{Im } \mathcal{A} - \alpha_0(S^1))$ the complement to $\text{Im } \mathcal{B} - \beta_0(S^1)$.

We claim that $\Psi^{-1} \circ g(y) = q_0 \circ \Psi^{-1}(y)$ for every point $y \in S^2$ in the complement to $\text{Im } \mathcal{B} - \beta_0(S^1)$. Indeed, the point $x = \Psi^{-1}(y)$ is well-defined and lies in $S^2 - j_0(\text{Im } \mathcal{A} - \alpha_0(S^1))$. The point $Q_0(x)$ cannot be in $\text{Im } \mathcal{A}$, otherwise

$$x \in Q_0^{-1}(\text{Im } \mathcal{A}) = j_0(f^{-1}(\text{Im } \mathcal{A})) \in j_0(\text{Im } \mathcal{A} - \alpha_0(S^1)).$$

Therefore, $j_0 \circ Q_0(x) = q_0(x) = q_0 \circ \Psi^{-1}(y)$ is well-defined. Clearly, $\Psi^{-1} \circ g(y)$ is also well-defined. Thus both parts of the equality $\Psi^{-1} \circ g = q_0 \circ \Psi^{-1}$ are defined on the complement to $\text{Im } \mathcal{B} - \beta_0(S^1)$. The equality itself holds wherever both parts are defined, because at least on some dense set, we have

$$\begin{aligned} \Psi^{-1} \circ g &= j_0 \circ \Phi^{-1} \circ \Phi \circ f \circ \Phi^{-1} = j_0 \circ f \circ \Phi^{-1} \\ &= j_0 \circ Q_0 \circ j_0 \circ \Phi^{-1} = q_0 \circ \Psi^{-1}. \end{aligned}$$

Note that the set $P_g \cup C_g$ is disjoint from $\text{Im } \mathcal{B} - \beta_0(S^1)$. Consider a simple curve Z disjoint from $\text{Im } \mathcal{B} - \beta_0(S^1)$ and containing $P_g \cup C_g$. (The existence of such curve follows from a simple Baire category argument, see Sect. 6.2 for more detail.) Then we can define a homeomorphism $\hat{\Psi} : S^2 \rightarrow S^2$ such that $\hat{\Psi}^{-1} = \Psi^{-1}$ on Z and on $g(Z)$. Indeed, both curves Z and $g(Z)$ are disjoint from $\text{Im } \mathcal{B} - \beta_0(S^1)$, thus Ψ^{-1} is well-defined on the union of these curves, and acts as a homeomorphism. It is not hard to see that Ψ^{-1} can be extended inside every component of the complement to $Z \cup g(Z)$ homeomorphically. Doing this for all components, we obtain a homeomorphism $\hat{\Psi}^{-1}$.

The restriction of g to Z is the same as the restriction of $\hat{\Psi} \circ q_0 \circ \hat{\Psi}^{-1}$ to Z . Indeed, for every $x \in Z$, we have

$$\hat{\Psi} \circ q_0 \circ \hat{\Psi}^{-1}(x) = \hat{\Psi} \circ q_0 \circ \Psi^{-1}(x) = \hat{\Psi} \circ \Psi^{-1} \circ g(x) = g(x).$$

We can even find a continuous ‘‘approximation’’ \hat{q}_0 of q_0 such that the restriction of $\hat{\Psi} \circ \hat{q}_0 \circ \hat{\Psi}^{-1}$ to Z is still the same. Indeed, we must have $q_0 = \hat{q}_0$ on

the set $\Psi^{-1}(Z)$, where q_0 is well-defined (as we know, q_0 is defined on all points $\Psi^{-1}(x)$, where $x \notin \text{Im } \mathcal{B} - \beta_0(S^1)$).

By Theorem 11, g is Thurston equivalent to $\hat{\Psi} \circ \hat{q}_0 \circ \hat{\Psi}^{-1}$, hence to \hat{q}_0 . \square

4.3 Semi-conjugacy

Recall the following theorem of Mary Rees [13]:

Theorem 12 *Suppose that a Thurston map F of degree 2 is Thurston equivalent to a hyperbolic rational function G . Moreover, suppose that there is an F -invariant complex structure near the critical orbits of F . Then there is a continuous map $\phi : S^2 \rightarrow S^2$ such that $\phi \circ F = G \circ \phi$.*

Proof We can assume F to be defined on $\overline{\mathbb{C}}$ (i.e. on a sphere with a global complex structure) and holomorphic on some open set U containing the post-critical set and satisfying $F(U) \Subset U$. We have the diagram

$$\begin{array}{ccc} \overline{\mathbb{C}} & \xrightarrow{F} & \overline{\mathbb{C}} \\ \downarrow \phi_1 & & \downarrow \phi_0 \\ \overline{\mathbb{C}} & \xrightarrow{G} & \overline{\mathbb{C}} \end{array}$$

where ϕ_0 and ϕ_1 are homeomorphisms holomorphic on U ; moreover, $\phi_1 = \phi_0$ on U , and ϕ_0 is isotopic to ϕ_1 relative to U (the existence of such homeomorphisms ϕ_0 and ϕ_1 follows from Böttcher’s theorem). Consider the multivalued function $G^{-1} \circ \phi_1 \circ F$. Since the critical values of $\phi_1 \circ F$ coincide with ramification points of G^{-1} , this multivalued function has a single valued branch ϕ_2 such that $\phi_2 = \phi_1$ on $F^{-1}(U)$. We have the following diagram:

$$\begin{array}{ccccc} \overline{\mathbb{C}} & \xrightarrow{F} & \overline{\mathbb{C}} & \xrightarrow{F} & \overline{\mathbb{C}} \\ \downarrow \phi_2 & & \downarrow \phi_1 & & \downarrow \phi_0 \\ \overline{\mathbb{C}} & \xrightarrow{G} & \overline{\mathbb{C}} & \xrightarrow{G} & \overline{\mathbb{C}} \end{array}$$

Similarly, we can define a sequence of homeomorphisms ϕ_n with the following properties: $G \circ \phi_n = \phi_{n-1} \circ F$ and $\phi_n = \phi_{n-1}$ on $F^{-(n-1)}(U)$. Then ϕ_n is a single valued branch of $G^{-n} \circ \phi_0 \circ F^{on}$.

We would like to prove that the sequence of maps ϕ_n converges uniformly. This would follow from the estimate

$$d(\phi_{n+1}(x), \phi_n(x)) \leq Cq^{-n},$$

where $0 < q < 1$ is a number independent of x . We can assume that $x \notin F^{-n}(U)$, otherwise the left-hand side is zero. Consider a curve γ connecting $\phi_0 \circ F^{on}(x)$ with $\phi_1 \circ F^{on}(x)$ in $\overline{\mathbb{C}} - \overline{\phi_0(U)}$ (we can arrange that this open set be connected by choosing a smaller U if necessary). Since $\overline{\mathbb{C}} - \overline{U}$ is compactly contained in $\overline{\mathbb{C}} - \overline{G \circ \phi_0(U)}$, the hyperbolic length of γ in $\overline{\mathbb{C}} - \overline{G \circ \phi_0(U)}$ can be made bounded by some constant independent of x and n . The length of the pull-back of γ under G^{on} is bounded by Cq^{-n} with $0 < q < 1$ by the Poincaré distance argument. The desired estimate now follows.

Let ϕ denote the limit of ϕ_n . Passing to the limit in both sides of the equality $G \circ \phi_n = \phi_{n-1} \circ F$, we obtain that $G \circ \phi = \phi \circ F$. □

We also need the following general fact:

Proposition 11 *Suppose that a continuous map $\phi : S^2 \rightarrow S^2$ is the limit of a uniformly convergent sequence of homeomorphisms ϕ_n . Then ϕ is onto and, for any point $v \in S^2$, the fiber $\phi^{-1}(v)$ is connected.*

Proof Any point y of $S^2 - \phi(S^2)$ has a neighborhood V such that $V \subset S^2 - \phi(S^2)$. We have $y = \phi_n(x_n)$ for all n . Since ϕ_n is uniformly close to ϕ for large n , we must have $\phi(x_n) \in V$, a contradiction.

Consider two points z and w in the fiber $\phi^{-1}(v)$. For large n , the points $\phi_n(z)$ and $\phi_n(w)$ are very close to each other. Let D_n be a small closed disk containing both of them, and set $A_n = \phi_n^{-1}(D_n)$. We can assume that the diameter of D_n tends to 0 as $n \rightarrow \infty$. The sequence of compact sets A_n has a subsequence that converges in the Hausdorff metric. Denote the limit by A . As a Hausdorff limit of compact connected sets, the set A is connected. Moreover, it contains both points z and w . We claim that $\phi(A) = v$. Indeed, for any point $a \in A$, there is a sequence $a_n \in A_n$ such that $a_n \rightarrow a$. The distance between $\phi_n(a_n)$ and $\phi(a)$ tends to zero, and $\phi_n(a_n) \rightarrow v$, hence $\phi(a) = v$. Thus any pair of points in $\phi^{-1}(v)$ belongs to a common connected subset of $\phi^{-1}(v)$. This means that $\phi^{-1}(v)$ is connected. □

Thus, in our setting, we obtain the following

Theorem 13 *The map h is semi-conjugate to g , i.e. there is a continuous surjective map $\phi : S^2 \rightarrow \overline{\mathbb{C}}$ such that $h \circ \phi = \phi \circ g$. Moreover, the fibers of ϕ are connected (i.e. ϕ is monotone).*

4.4 Dynamics of g

In this section, we discuss some simple dynamical properties of g that we need for the proof that g is conjugate to h .

Lemma 3 *There is a positive real number δ_0 such that for every pair $x, y \in J_g$ with $0 < d(x, y) < \delta_0$, we have $g(x) \neq g(y)$.*

Proof Since there are no critical points in J_g , every point $x \in J_g$ has a neighborhood U_x such that $g|_{U_x}$ is injective. Let δ_0 be the Lebesgue number of the covering $\mathcal{U} = \{U_x\}_{x \in J_g}$. Now assume that $0 < d(x, y) < \delta_0$. Then $x, y \in U$, where $U \in \mathcal{U}$. Since g is injective on U , the result follows. □

Lemma 4 *There is a self-homeomorphism $M \neq id$ of the dynamical sphere of g such that $M^{\circ 2} = id$, and $g \circ M = g$. Moreover, we have $\phi \circ M = -\phi$, where ϕ is the map from Theorem 13.*

Proof Indeed, the multivalued map $g^{-1} \circ g$ splits into two single valued branches. One of these branches is the identity transformation. Let M be the other branch. We have $g \circ M = g$ by definition. It follows that $g \circ M^{\circ 2} = g$, thus $M^{\circ 2}$ is either M or the identity. Since $M \neq id$, we have $M^{\circ 2} \neq M$. Therefore, $M^{\circ 2} = id$.

We have

$$h \circ \phi \circ M = \phi \circ g \circ M = \phi \circ g = h \circ \phi,$$

therefore $\phi(M(x)) = \pm\phi(x)$ for all x in the dynamical sphere of g . Near the critical points of g (which are the ϕ -preimages of 0 and ∞), the map ϕ is a homeomorphism, and we have the minus sign. By continuity, the sign is the same for all points. □

4.5 Triviality of fibers

In this section, we prove that the continuous map ϕ from Theorem 13 is actually a homeomorphism. Note first that the restriction of ϕ to the Fatou set of g is injective. This follows from the definition of ϕ given in the proof of Theorem 12. Indeed, on every compact set contained in the Fatou set of g , the map ϕ coincides with a homeomorphism ϕ_n . We know that fibers of ϕ are connected. Thus they lie entirely in J_g (a connected set intersecting the Fatou set of g by just one point is itself a singleton).

Lemma 5 *The restriction of g to any fiber of ϕ is injective.*

Proof Assume the contrary: $g(x) = g(y)$ for a pair of different points x, y such that $\phi(x) = \phi(y)$. Let M be the involution introduced in Lemma 4. Then $y = M(x)$, and

$$\phi(y) = \phi(M(x)) = -\phi(x) = -\phi(y).$$

It follows that $\phi(y) = 0$ or ∞ , a contradiction. □

Lemma 6 *The image of any fiber of ϕ under the map g is contained in a fiber of ϕ .*

Proof Indeed, let Z be a fiber of ϕ . We have $\phi(g(Z)) = h(\phi(Z))$, which is a single point. □

The following proposition is a minor modification of an argument given in [13].

Proposition 12 *Suppose that ϕ is not a homeomorphism, i.e. it has a non-trivial fiber. Let δ_0 be the number introduced in Lemma 3. For every $\varepsilon > 0$, there exist $x, y \in J_g$ such that*

$$d(x, y) > \delta_0/2, \quad d(g(x), g(y)) < \varepsilon, \quad \phi(x) = \phi(y).$$

Proof Without loss of generality, we can assume that $\varepsilon < \delta_0/2$. Let N be a positive integer such that every subset of J_g with at least N points has a pair of distinct points on distance $< \delta(\varepsilon)$ (where $\delta(\varepsilon)$ is as in Theorem 9). Consider a nontrivial fiber Z of ϕ , and a subset Z_0 of Z containing N points. (As a connected set of cardinality > 1 , the set Z must actually have uncountably many points.) Define ε' to be the minimal distance between different points in Z_0 . Set $n_0 = n_0(\varepsilon, \varepsilon')$ (where $n_0(\varepsilon, \varepsilon')$ is as in Theorem 9). The set $g^{on_0}(Z_0)$ has cardinality N . This follows from Lemmas 5 and 6. Therefore, there is a pair of points x_{n_0}, y_{n_0} in this set such that $d(x_{n_0}, y_{n_0}) < \delta(\varepsilon)$. Note that $\delta(\varepsilon) \leq \varepsilon$, in particular, $d(x_{n_0}, y_{n_0}) < \varepsilon$.

For every $k = 0, \dots, n_0$, we can define x_k and y_k inductively by the following relations:

$$x_k, y_k \in g^{ok}(Z_0), \quad g(x_k) = x_{k+1}, \quad g(y_k) = y_{k+1}.$$

Then either y_k is the closest to x_k preimage of y_{k+1} and, by Proposition 10, we have $d(x_k, y_k) < \varepsilon$, or $d(x_k, y_k) \geq \delta_0 - \varepsilon > \delta_0/2$. Indeed, let y'_k be the closest to x_k preimage of y_{k+1} . If $y'_k \neq y_k$, then

$$d(x_k, y_k) \geq d(y_k, y'_k) - d(y'_k, x_k) \geq \delta_0 - \varepsilon.$$

Suppose first that $d(x_k, y_k) \geq \delta_0/2$ for some k . Then $x = x_k$ and $y = y_k$ have the required properties. Suppose now that y_k is the closest to x_k preimage of y_{k+1} , for all k . Then we have $d(x_0, y_0) < \varepsilon'$ by Proposition 10. But this contradicts our choice of ε' , since $x_0, y_0 \in Z_0$. □

Suppose that ϕ is not a homeomorphism. Then, by Proposition 12, there exist points x_n and y_n in J_g such that

$$d(x_n, y_n) \geq \delta_0/2, \quad d(g(x_n), g(y_n)) < 1/n, \quad \phi(x_n) = \phi(y_n).$$

Passing to suitable subsequences if necessary, we can assume that x_n and y_n converge to some points x and y in J_g , respectively. For these limit points, we must have

$$d(x, y) \geq \delta_0/2, \quad g(x) = g(y), \quad \phi(x) = \phi(y).$$

In other terms, x and y belong to the same fiber of ϕ , and the restriction of g to this fiber is not injective. This contradicts Lemma 5. The contradiction shows that all fibers of ϕ are trivial, thus ϕ is a homeomorphism.

This finishes the proof of Theorems 8 and 1.

5 Holomorphic regluing

In this section, we just sketch main characters of holomorphic theory of regluing, and give the most basic constructions.

5.1 An example

We consider first a simple example, where we define an explicit sequence of approximations to a regluing. This sequence will consist of partially defined but holomorphic functions.

Let f be the quadratic polynomial $z \mapsto z^2 - 6$. The Julia set J_f of f is a Cantor set that lies on the real line. Recall that the biggest component of the complement to J_f in $[-3, 3]$ is $(-\sqrt{3}, \sqrt{3})$. Suppose we want to reglue the interval $[-\sqrt{3}, \sqrt{3}]$, thus connecting two parts of J_f . This is done by the following map:

$$j(z) = \sqrt{z^2 - 3}$$

(which is understood as a branch over the complement to $[-\sqrt{3}, \sqrt{3}]$ that is tangent to the identity at infinity). The inverse map is given by the formula $j^{-1}(z) = \sqrt{z^2 + 3}$, and is defined on the complement to $[-i\sqrt{3}, i\sqrt{3}]$.

Consider the composition $f \circ j^{-1}$. It turns out that this function extends to a quadratic polynomial! Indeed, we have

$$f(\sqrt{z^2 + 3}) = (z^2 + 3) - 6 = z^2 - 3.$$

We denote the polynomial in the right-hand side by p_{-3} ; in general, p_c stands for the quadratic polynomial $z \mapsto z^2 + c$. On a more conceptual level, in order to see that $f \circ j^{-1}$ is a restriction of a polynomial, it suffices to show that it extends continuously to the interval $[-i\sqrt{3}, i\sqrt{3}]$. This follows from the fact that f folds the interval $[-\sqrt{3}, \sqrt{3}]$ at 0.

In fact, what we really want to consider is not the composition but the “conjugation” $j \circ f \circ j^{-1}$. Since $f \circ j^{-1}$ is a restriction of the polynomial p_{-3} ,

we define the new function f_1 as $j \circ p_{-3}$, which has a larger domain than $j \circ f \circ j^{-1}$. Unfortunately, the function f_1 is not continuous. The discontinuity of this function is due to the discontinuity of j . Actually, the function f_1 is defined and holomorphic on the complement to two simple curves—the pullbacks of $[-\sqrt{3}, \sqrt{3}]$ under p_{-3} , and it “reglues” these curves in a sense. We would like to get rid of this discontinuity by “conjugating” f_1 with yet another regluing map. To this end, we need an injective holomorphic function j_1 defined on the domain of f_1 and having the same type of discontinuity at the two special curves, where f_1 is undefined. We cannot take $j_1 = f_1$ because f_1 is, in general, two-to-one (it is the square root of a degree four polynomial). However, we can take

$$j_1 = \sqrt{f_1 - f_1(0)}.$$

The square root may look disturbing but it does not actually create any ramification, so that the function j_1 is a union of two branches. These branches are still not everywhere defined (they are defined exactly on the domain of f_1) but they are single valued! Indeed, the square root has ramification points exactly where f_1 is equal to $f_1(0)$ or ∞ . But at all such places, namely at 0 and at ∞ , the function $\sqrt{f_1}$ has simple critical points. Thus at these places, the function j_1 looks like $\sqrt{u^2}$, where u is some local coordinate, and this does not have any ramification. It is also easy to see that j_1 has no nontrivial monodromy around the curves, on which it is undefined (it suffices to look at the monodromy of the square root). We choose the branch of j_1 that is tangent to the identity at infinity.

Now finding $f_1 \circ j_1^{-1}$ is easy: Set $w = j_1(z)$, then $w = \sqrt{f_1(z) - f_1(0)}$, and

$$f_1 \circ j_1^{-1}(w) = f_1(z) = w^2 + f_1(0)$$

for all w such that the left-hand side is defined. So this is again a quadratic polynomial! By the way, the number $c_1 = f_1(0)$ is easy to compute:

$$f_1(0) = j(-3) = -\sqrt{6} = -2^{1/2} \cdot 3^{1/2}.$$

The only non-trivial part is the sign of the square root. It is determined by our choice of the branch for j but we skip the corresponding computation.

Next, we define the function $f_2 = j_1 \circ p_{c_1}$. Note that $j_1 = p_{c_1}^{-1} \circ f_1$ on the domain of f_1 . It follows that

$$f_2 \text{ “=” } p_{c_1}^{-1} \circ f_1 \circ p_{c_1}.$$

This formula looks nice but one needs to be very careful, because the expression in the right-hand side is ambiguous (it should be considered as a single-valued branch over some domain, but what is written does not carry

any information on the domain of definition). The right understanding of this formula is that $p_{c_1}^{-1} \circ f_1$ should be thought of as j_1 but then it coincides with the formula $f_2 = j_1 \circ p_{c_1}$ that we have used to define f_2 . In our formulas, one can recognize Thurston’s iteration but in a slightly unusual context because we deal with discontinuous holomorphic functions rather than with continuous non-holomorphic functions. There is a precise relation between what is happening and Thurston’s theory (better seen on other examples, because, in the case under consideration, the Teichmüller space has dimension 0, and Thurston’s theory does not have much to say).

Continuing the same process, we obtain a sequence f_n of functions, each defined on the complement to a finite union of simple curves, with the following recurrence property:

$$f_{n+1} = j_n^{-1} \circ p_{c_n}, \quad c_n = f_n(0),$$

where j_n is a branch of $\sqrt{f_n - c_n}$ defined on the domain of f_n and tangent to the identity at infinity.

In our example, we can compute the numbers c_n explicitly. Consider the sequence $f_1^{om}(0)$. This sequence stabilizes at the second term:

$$-\sqrt{6}, \quad \sqrt{6}, \quad \sqrt{6}, \quad \dots, \quad \sqrt{6}, \quad \dots$$

The sequence $f_2^{om}(0)$ can be obtained from the first sequence as follows:

$$f_2^{om}(0) = \sqrt{f_1^{om+1}(0) - f_1(0)}.$$

Thus all terms are equal to $\pm\sqrt{2\sqrt{6}} = \pm 2^{1-1/4} \cdot 3^{1/4}$. The determination of signs is a bit tricky, but the correct signs are the following: the first sign is minus, and all other signs are plus. Continuing in the same way, we obtain that

$$f_n^{om}(0) = \pm 2^{1-1/2^n} \cdot 3^{1/2^n},$$

where the first sign is minus, and all other signs are plus. In particular, $c_n = f_n(0) \rightarrow -2$ as $n \rightarrow \infty$, and the convergence is exponentially fast.

We see that the sequence of polynomials p_{c_n} converges to the polynomial $g : z \mapsto z^2 - 2$. This is the so called *Tchebyshev polynomial*. The Julia set of this polynomial is equal to the interval $[-2, 2]$. Note that the orbit of the critical point 0 is finite: $0 \mapsto -2 \mapsto 2$, and 2 is fixed.

Define the sequence of maps $\Phi_n = j_n \circ \dots \circ j_1 \circ j$. The map Φ_n is defined and holomorphic on the complement to finitely many simple curves. The main property of Φ_n is that $f_{n+1} = \Phi_n \circ f \circ \Phi_n^{-1}$ wherever the right-hand side is defined, which follows from the definition. Note also that $p_{c_n} = \Phi_{n-1} \circ f \circ \Phi_{n-1}^{-1}$ wherever the right-hand side is defined. In our example, it can be

shown that the sequence Φ_n converges uniformly to a map Φ defined on the complement to countably many curves—in our case, to all iterated pullbacks of $[-\sqrt{3}, \sqrt{3}]$ under f . The map Φ reglues all these horizontal intervals into “vertical curves” so that the Julia set of f , which is a Cantor set, gets glued into the interval $[-2, 2]$ under Φ . Passing to the limit as $n \rightarrow \infty$ in the identity $p_{c_n} = \Phi_{n-1} \circ f \circ \Phi_n^{-1}$, we obtain that

$$g = \Phi \circ f \circ \Phi^{-1}.$$

Note that the right-hand side is only defined on the complement to countably many curves, but it extends to the complex plane as a holomorphic function.

5.2 Thurston’s algorithm for quadratic branched coverings

In the example above, we constructed an explicit sequence of approximations to a topological regluing. We will now define these approximations in a more general context. First, we recall a version of the Thurston algorithm for branched coverings of degree two, and then extend this algorithm to certain classes of partially defined maps.

Lemma 7 *For any pair of different points $a, b \in \overline{\mathbb{C}}$, there exists a quadratic rational function with critical values a and b .*

An analog of this statement for arbitrary degree is true and well known but the quadratic case is more explicit.

Proof Set

$$R_{a,b}(z) = \begin{cases} \frac{az^2-b}{z^2-1}, & a, b \neq \infty, \\ z^2 + b, & a = \infty, b \in \mathbb{C}, \\ z^{-2} + a, & a \in \mathbb{C}, b = \infty. \end{cases}$$

Then the critical points of $R_{a,b}$ are ∞ and 0 , and we have

$$R_{a,b}(\infty) = a, \quad R_{a,b}(0) = b. \quad \square$$

Suppose we start with a branched covering $f_1 : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$. Let R_1 be a rational function, whose critical values coincide with those of f_1 . Then there is a global branch j_1 of the multi-valued function $R_1^{-1} \circ f_1$. Note that R_1 is only defined up to pre-composition with a Möbius transformation. Hence j_1 is only defined up to post-composition with a Möbius transformation. Consider the map $f_2 = j_1 \circ R_1$. This map is an extension of $j_1 \circ f \circ j_1^{-1}$, hence it is only defined up to a Möbius conjugacy. Continue the same process with

function f_2 : consider a quadratic rational function R_2 with the same critical values, define j_2 as a branch of $R_2^{-1} \circ f_2$, etc. We obtain the following diagram:

$$\begin{array}{ccccccc}
 \overline{\mathbb{C}} & \xrightarrow{j_1} & \overline{\mathbb{C}} & \xrightarrow{j_2} & \overline{\mathbb{C}} & \xrightarrow{j_3} & \overline{\mathbb{C}} \longrightarrow \dots \\
 f_1 \downarrow & & f_2 \downarrow & & f_3 \downarrow & & f_4 \downarrow \\
 \overline{\mathbb{C}} & \xrightarrow{j_1} & \overline{\mathbb{C}} & \xrightarrow{j_2} & \overline{\mathbb{C}} & \xrightarrow{j_3} & \overline{\mathbb{C}} \longrightarrow \dots
 \end{array}$$

The maps j_n are only defined up to post-composition with Möbius transformations. Therefore, we need to make some normalization in order to fix these maps. Below, a natural normalization is suggested for quadratic rational functions. We can ask then the following questions:

1. does the sequence of rational functions R_n converge?
2. does the sequence of maps $\Psi_n = j_n \circ \dots \circ j_1$ converge?

Convergence is understood as uniform convergence with respect to the spherical metric.

It is not hard to show that if Ψ_n converge to a homeomorphism Ψ , then the rational functions

$$R_n = \Psi_{n-1} \circ f_1 \circ \Psi_n^{-1}$$

converge to a rational function R_∞ . Moreover, Ψ conjugates f_1 with R_∞ . If Ψ_n converge to a function Ψ (not necessarily a homeomorphism), and the sequence R_n is equicontinuous, then it also converges. Indeed, for every $\varepsilon > 0$, the uniform distance between Ψ_n and Ψ_m is less than ε for sufficiently large n and m . Since all Ψ_n are surjective, it follows that the uniform distance between $j_m \circ \dots \circ j_{n+1}$ and the identity is less than ε . Finally, we can conclude that R_n and

$$R_m = (j_{m-1} \circ \dots \circ j_n) \circ R_n \circ (j_m \circ \dots \circ j_{n+1})^{-1}$$

are uniformly close. Thus R_n converge to some rational function R_∞ , and Ψ semi-conjugates f_1 with R_∞ .

Suppose now that f_1 is critically finite. Then Thurston’s theory provides a technique to answer the first question. The answer to the second question depends on a particular normalization for j_n . However, once the first question is answered in the affirmative, it is in general much easier to answer the second question.

5.3 A modification for partially defined maps

We need to consider the case, where f_1 is only defined on a part of the sphere (e.g. on the complement to finitely many simple curves). Thurston’s algorithm (as described above) can be modified in the following way. Assume that f_1 is defined on some open subset $U_1 \subset S^2$, and that $f_1 : U_1 \rightarrow f_1(U_1)$ is an orientation-preserving branched covering of degree two with two critical points. Moreover, assume that forward f_1 -orbits of both critical points are well-defined (i.e. lie in U_1). Then we can proceed as before: choose a quadratic rational function R_1 , whose critical values coincide with those of f_1 . It is not hard to show that the multi-valued function $R_1^{-1} \circ f_1$ has trivial monodromy, hence splits into two single valued branches.

Let j_1 be one of the branches. It is defined on the set U_1 . Form a new function $f_2 = j_1 \circ R_1$. It is defined on the set $R_1^{-1}(U_1)$. We need to prove that the critical orbits of f_2 are well-defined. Indeed, let c_1 and c_2 be the critical points of f_1 . We know that the points $f_1^{on}(c_i)$ lie in U_1 for all $n \geq 0$ and $i = 1, 2$. Therefore, the points $j_1 \circ f_1^{on}(c_i)$ are well-defined. Note also that $j_1(c_i)$ are the critical points of f_2 , and that

$$j_1 \circ f_1^{on}(c_i) = f_2^{on} \circ j_1(c_i).$$

(We use repeatedly that $f_2 = j_1 \circ R_1$ and $f_1 = R_1 \circ j_1$.) In particular, the right-hand side is defined. We can now repeat the step of Thurston’s algorithm. In this way, we obtain a sequence of maps f_n , j_n and rational functions R_n , as above. The only difference is that f_n and j_n are not everywhere defined.

We can now be more specific regarding the normalization of j_n . Suppose that the critical points of f_1 are ∞ and 0 (this can be always arranged by a suitable Möbius conjugacy). Let $a_1 = f_1(\infty)$ and $b_1 = f_1(0)$ be the corresponding critical values. We choose R_1 in the form R_{a_1, b_1} , where the rational functions $R_{a, b}$ were introduced in Lemma 7. This fixes R_1 , however, we still need to decide between the two branches of $R_1^{-1} \circ f_1$. We will do this in two examples. These examples are, however, the most important.

Example 2 Suppose that $a_1 = f_1(\infty) = \infty$, and that $f_1(z) = z^2 + o(z^2)$ near infinity. Then both branches of $R_1^{-1} \circ f_1$ fix ∞ . However, only one branch is tangent to the identity at infinity. We choose j_1 to be this branch. Note by the way that $j_1(z) = R_1^{-1} \circ f_1(z)$ is given by the following explicit formula

$$\sqrt{f_1(z) - b_1}.$$

We have $f_2(z) = z^2 + o(z^2)$ near ∞ so that we can choose the branch for j_2 in the same way, etc.

Example 3 Suppose that $f_1(1) = \infty$ and that ∞ is not a critical value of f_1 . Then one branch of $R_1^{-1} \circ f_1$ maps 1 to 1, and the other branch to -1 . Indeed, $R_1(1) = R_1(-1) = \infty = f_1(1)$. We choose j_1 to be the branch that fixes 1. Note that $j_1(z) = R_1^{-1} \circ f_1(z)$ is given by the following explicit formula

$$\sqrt{\frac{f_1(z) - b_1}{f_1(z) - a_1}}$$

By definition, j_1 fixes both 0 and ∞ . Therefore, $f_2 = j_1 \circ f_1$ also maps 1 to ∞ , and ∞ is not a critical value of f_2 . Thus we can choose the branch for j_2 in the same way, etc.

5.4 The first regluing

Let us start with a quadratic rational function f . Consider a simple path $\alpha_0 : [-1, 1] \rightarrow \overline{\mathbb{C}}$ such that $f \circ \alpha_0(-t) = f \circ \alpha_0(t)$ for all $t \in [-1, 1]$. In particular, $\alpha_0(0)$ must be a critical point of f . The path α_0 can be interpreted as the α -path

$$(t_1, t_2) \mapsto \alpha_0(t_1).$$

We will sometimes use this interpretation when referring to regluing, e.g. regluing of α_0 means regluing of this α -path. To fix ideas, we assume that the critical points of f are 0 and ∞ and that $\alpha_0(0) = 0$. If all pullbacks of α_0 are disjoint, then we can consider a topological regluing of these pullbacks, as defined earlier in the text. However, we will now make a much weaker assumption, namely, that the orbits of $f \circ \alpha_0(1)$ and ∞ are disjoint from $\alpha_0[-1, 1]$.

Let R be a rational function, whose critical values are $f \circ \alpha_0(1)$ and $f(\infty)$. (E.g. we can choose R to have the form $R_{a,b}$ as in Lemma 7.) Consider the multi-valued analytic function $R^{-1} \circ f$. There are two single valued branches of this function over the complement to $\alpha_0[-1, 1]$. We choose one branch, and call it j . Now $f_1 = j \circ R$ satisfies the assumptions of Sect. 5.3, and we can run Thurston’s algorithm for f_1 . In the following examples, we comment on a specific choice of the branch for j .

Example 4 Suppose that f is a quadratic polynomial. In a suitable coordinate, it can be written in the form $f(z) = z^2 + b$. The rational function R should have critical values $\alpha_0(1)^2 + b$ and ∞ , thus we can take $R(z) = z^2 + \alpha_0(1)^2 + b$. The multi-valued function $R^{-1} \circ f$ is given by the following explicit formula:

$$z \mapsto \sqrt{z^2 - \alpha_0(1)^2}.$$

This function has two branches over the complement to $\alpha_0[-1, 1]$. One of the branches is tangent to the identity near infinity. We choose j to be this branch. For the corresponding function $f_1 = j \circ R$, we have $f_1 = z^2 + o(z^2)$ near infinity. Therefore, the normalization of j_n can be made as in Example 2.

Example 5 Consider a quadratic rational function f that is conjugate neither to a polynomial nor to the map $z \mapsto 1/z^2$. Then we can reduce f to the form

$$z \mapsto \frac{az^2 - b}{z^2 - 1}$$

by a suitable Möbius conjugacy. Note that $a = f(\infty)$ and $b = f(0)$. The rational function R should have critical values $f \circ \alpha_0(1)$ and $f(\infty) = a$. Therefore, we can take

$$R(z) = \frac{az^2 - f \circ \alpha_0(1)}{z^2 - 1}.$$

The multi-valued function $R^{-1} \circ f$ is given by the following explicit formula:

$$z \mapsto \sqrt{\frac{z^2 - \alpha_0(1)^2}{1 - \alpha_0(1)^2}}.$$

Note that one branch of this function takes 1 to 1, and the other branch to -1 . We define j as the branch that fixes 1. The function $f_1 = j \circ R$ takes 1 to ∞ , and ∞ is not a critical value of f_1 . Therefore, we can use normalization for j_n introduced in Example 3.

5.5 Some questions

Starting with a quadratic rational function f and a simple path $\alpha_0 : [-1, 1] \rightarrow \overline{\mathbb{C}}$, we have defined sequences of maps f_n, j_n and rational functions R_n (provided that α_0 satisfies certain properties given in Sect. 5.4). For every n , the functions f_n and j_n are defined on a complement to finitely many simple curves (possibly intersecting). As before, we can define the maps $\Psi_n = j_n \circ \dots \circ j_1$. The map Ψ_n is defined on a set U_n , which is also a complement to finitely many simple curves. Since every U_n is open and dense, the intersection $\bigcap U_n$ is dense. We can ask whether the maps Ψ_n converge uniformly on $\bigcap U_n$.

In the case where f_1 is critically finite, this question is closely related to Thurston’s theory. This theory gives an answer to the question on convergence of R_n .

Suppose that R_n and Ψ_n converge uniformly. Then the limit Ψ of Ψ_n is a partially defined semi-conjugacy between f_1 and the rational function

$R_\infty = \lim R_n$ (meaning that Ψ semi-conjugates the restriction of f_1 to a certain forward invariant dense subset with the restriction of R_∞ to a certain forward invariant dense subset). Moreover, $\Phi = \Psi \circ j$ is a partially defined semi-conjugacy between f and R_∞ .

What can be said about the limit map Ψ ? To what extent is this map holomorphic? Note that it is holomorphic in the interior of $\bigcap U_n$ (which may well be empty) but the notion of being holomorphic is not even defined on the boundary. Is it possible to define an appropriate notion generalizing holomorphy to the boundary points so that Ψ would be holomorphic everywhere on $\bigcap U_n$? (An attempt to define this notion is made in preprint [18].)

6 Existence of topological regluing

In this section, we prove Theorems 3 and 4. Our methods are based on a theory of Moore [10] that gives a purely topological characterization of topological spheres.

6.1 A variant of Moore's theory

Moore [10] defined a system of topological conditions that are necessary and sufficient for a topological space to be homeomorphic to the sphere. He used this system to lay axiomatic foundations of plane topology. One of the most remarkable applications of Moore's theory is a description of equivalence relations on the sphere such that the quotient space is homeomorphic to the sphere. Moore's theory has been further developed in [2, 20, 21], see also [3].

In this section, we will give a "rapid introduction" into a version of Moore's theory. Our axiomatics is very far from being optimal (many axioms can be proved as theorems), and it does not have the purpose of setting a foundation to plane topology, but just serves as a fast working tool to prove that something is a topological sphere. A better system of axioms is given in [19].

Let X be a compact connected Hausdorff space. Recall that a *simple closed curve* in X is the image in X of a continuous embedding $\gamma : S^1 \rightarrow X$. Here the map γ is called a *simple closed path*. We also define a *simple path* as a continuous embedding of $[0, 1]$ into X , and a *simple curve* as the image of a simple path. A *segment* of a simple curve is defined as the image of a closed subinterval in $[0, 1]$ under the corresponding simple path. Similarly, we can define a segment (or an arc) of a simple closed curve.

Suppose we fixed some set \mathcal{E} of simple curves in X . The curves in \mathcal{E} are called *elementary curves*. We will always assume that segments of elementary curves are also elementary curves, and that if two elementary curves have only an endpoint in common, then their union is also an elementary curve. We will not state these assumptions as axioms, although, technically, they are. Define

an *elementary closed curve* as a simple closed curve, all of whose segments are elementary curves.

We are ready to state the first axiom:

Axiom 1 (Elementary domain axiom) Any elementary closed curve J divides X into two connected components called *elementary domains* bounded by J .

Since a simple closed curve is homeomorphic to a circle, it makes sense to talk about the (circular) order of points in it. Define an *elementary quadrilateral* in X as an elementary domain bounded by an elementary closed curve $J \subset X$ with a distinguished quadruple of different points a, b, c and d in J . We assume that the points a, b, c and d appear in J in the same circular order as they are listed. Consider an elementary quadrilateral Q bounded by J . A simple curve connecting the segment $[a, b]$ with the segment $[c, d]$ of J is called a *vertical curve*, provided that it lies entirely in Q , except for the endpoints. A simple curve connecting the segments $[b, c]$ and $[d, a]$ is called *horizontal*, provided that it lies entirely in Q , except for the endpoints. We will also regard $[a, b]$ and $[c, d]$ as horizontal curves, $[b, c]$ and $[d, a]$ as vertical curves.

Axiom 2 (Extension axiom) For any elementary quadrilateral Q bounded by an elementary closed curve J , and any point x in J , there exists a vertical or a horizontal elementary curve with an endpoint at x . Moreover, this curve divides Q into at most two elementary quadrilaterals.

From the Extension axiom, it follows easily that the topological boundary of any elementary domain bounded by an elementary closed curve J coincides with J .

Define a *grid* in an elementary quadrilateral Q as a system of finitely many horizontal elementary curves and finitely many vertical elementary curves such that all horizontal curves are pair-wise disjoint, all vertical curves are pair-wise disjoint, and every horizontal curve intersects every vertical curve at exactly one point. Using the Elementary domain axiom and the Extension axiom, it is not hard to show that every grid with $n - 1$ horizontal and $m - 1$ vertical curves divides Q into mn pieces. We will refer to these pieces as *cells of the grid*. Cells can be regarded as elementary quadrilaterals.

Axiom 3 (Covering axiom) Consider an elementary quadrilateral Q and an open covering \mathcal{U} of \overline{Q} . Then there exists a grid in Q such that the closure of every cell lies in some element of \mathcal{U} (such grid is said to be *subordinate* to \mathcal{U}).

The following is the main topological fact we need.

Theorem 14 *Let X be a compact connected Hausdorff second countable topological space. Suppose that a set of elementary curves \mathcal{E} in X satisfies the Elementary domain axiom, the Extension axiom and the Covering axiom. Suppose also that there exists an elementary closed curve in X . Then X is homeomorphic to the sphere.*

Proof Since there exists an elementary closed curve, by the Elementary domain axiom, there exists an elementary quadrilateral Q . It suffices to prove that the closure of this quadrilateral is homeomorphic to the closed disk.

Fix a countable basis \mathcal{B} of the topology in X . There are countably many finite open coverings of \overline{Q} contained in \mathcal{B} . Number all such coverings by natural numbers. We will define a sequence of grids G_n in Q by induction on n . For $n = 1$, we just take the trivial grid, the one that does not have any horizontal or vertical curves. Suppose now that G_n is defined. Let \mathcal{U}_n be the n -th covering of \overline{Q} . Using the Covering axiom, we can find a grid in each cell of G_n that is subordinate to \mathcal{U}_n . Using the Extension axiom, we can extend these grids to a single grid G_{n+1} in Q . Thus G_{n+1} contains G_n and is subordinate to \mathcal{U}_n .

Consider any pair of different points $x, y \in \overline{Q}$. There exists n such that x and y do not belong to the closure of the same cell in G_{n+1} . Indeed, let us first define a covering of \overline{Q} as follows. For any $z \in \overline{Q}$, choose U_z to be an element of \mathcal{B} that contains z but does not include the set $\{x, y\}$. The sets U_z form an open covering of \overline{Q} . Since \overline{Q} is compact, there is a finite subcovering. This finite subcovering coincides with \mathcal{U}_n for some n . Then, by our construction, the closure of every cell in G_{n+1} is contained in an element of \mathcal{U}_n . However, the set $\{x, y\}$ is not contained in an element of \mathcal{U}_n . Therefore, $\{x, y\}$ cannot belong to the closure of a single cell.

Consider a nested sequence $C_1 \supset C_2 \supset \dots \supset C_n \supset \dots$, where C_n is a cell of G_n . We claim that the intersection of the closures \overline{C}_n is a single point. Indeed, this intersection is nonempty, since \overline{C}_n form a nested sequence of nonempty compact sets. On the other hand, as we saw, there is no pair $\{x, y\}$ of different points contained in all \overline{C}_n .

Consider the standard square $[0, 1] \times [0, 1]$ and a sequence H_n of grids in it with the following properties:

1. all horizontal curves in H_n are horizontal straight line segments, all vertical curves in H_n are vertical straight line segments;
2. the grid H_n has the same number of horizontal curves and the same number of vertical curves as G_n , thus there is a natural one-to-one correspondence between cells of H_n and cells of G_n respecting ‘‘combinatorics’’, i.e. the cells in H_n corresponding to adjacent cells in G_n are also adjacent;
3. the grid H_{n+1} contains H_n ; moreover, if a cell of H_{n+1} is in a cell of H_n , then there is a similar inclusion between the corresponding cells of G_{n+1} and G_n ;

4. between any horizontal curve of H_n and the next horizontal curve, the horizontal curves of H_{n+1} are equally spaced; similarly, between any vertical curve of H_n and the next vertical curve, the vertical curves of H_{n+1} are equally spaced.

It is not hard to see that any nested sequence of cells D_n of H_n converges to a point: $\bigcap \overline{D}_n = \{pt\}$.

We can now define a map $\Phi : \overline{Q} \rightarrow [0, 1] \times [0, 1]$ as follows. For a point $x \in \overline{Q}$, there is a nested sequence of cells C_n of G_n such that \overline{C}_n contains x for all n . Define the point $\Phi(x)$ as the intersection of the closures of the corresponding cells D_n in H_n . (Note that they also form a nested sequence according to our assumptions.) Clearly, the point $\Phi(x)$ does not depend on a particular choice of the nested sequence C_n (there can be at most four different choices). It is also easy to see that Φ is a homeomorphism between \overline{Q} and the standard square $[0, 1] \times [0, 1]$. □

One of the main applications of Moore’s theory is the following characterization of equivalence relations on S^2 such that the corresponding quotient spaces are homeomorphic to the sphere [11]:

Theorem 15 *Let \sim be a closed equivalence relation on S^2 such that all equivalence classes are connected and do not separate the sphere. Then the quotient S^2 / \sim is homeomorphic to the sphere provided that not all points are equivalent.*

Recall that a closed equivalence relation on S^2 is an equivalence relation represented by a closed subset of $S^2 \times S^2$.

6.2 Blow-up space

Consider a countable set \mathcal{Z} of disjoint compact connected locally connected nonseparating sets in S^2 . Suppose that \mathcal{Z} forms a null-sequence. For every $A \in \mathcal{Z}$, fix a continuous map $\Pi_A : S^2 \rightarrow S^2$ such that Π_A restricts to an orientation-preserving homeomorphism between Δ_∞ and $S^2 - A$, and $\Pi_A(\overline{\Delta}) = A$ (we use notation from Sect. 2.1).

We can now define the *blow-up space* $X_{\mathcal{Z}}$ of \mathcal{Z} as the equalizer of all maps Π_A , i.e. the subset of the product $(S^2)^{\mathcal{Z}}$ consisting of all functions $\chi : \mathcal{Z} \rightarrow S^2$ such that the points $\Pi_A(\chi(A)) \in S^2$ coincide for all $A \in \mathcal{Z}$. Intuitively, $X_{\mathcal{Z}}$ is obtained from S^2 by blowing up every $A \in \mathcal{Z}$ according to the map Π_A . As a closed subset in the compact Hausdorff second countable space $(S^2)^{\mathcal{Z}}$, the space $X_{\mathcal{Z}}$ is also compact, Hausdorff and second countable. As a continuous map from a compact space to a Hausdorff space, the map $\Pi_{\mathcal{Z}}$ is closed. For $A \in \mathcal{Z}$, define the subset $\Delta_A \subset X_{\mathcal{Z}}$ as the set of points $\chi \in X_{\mathcal{Z}}$ such that $\chi(A) \in \Delta$. Clearly, Δ_A is homeomorphic to the open disk Δ , and

$\overline{\Delta}_A$ is homeomorphic to the closed disk $\overline{\Delta}$. The topological disks $\overline{\Delta}_A, A \in \mathcal{Z}$, are disjoint and form a null-sequence. Intuitively, the sets $A \in \mathcal{Z}$ are blown up to disks $\overline{\Delta}_A$.

The main result of this section is the following

Theorem 16 *Under our assumptions on \mathcal{Z} , the blow up space $X_{\mathcal{Z}}$ is homeomorphic to the sphere.*

To prove Theorem 16, we will use the technique introduced in Sect. 6.1. First note that there is a natural projection $\Pi_{\mathcal{Z}} : X_{\mathcal{Z}} \rightarrow S^2$ that takes $\chi \in X_{\mathcal{Z}}$ to the point $\Pi_A(\chi(A)), A \in \mathcal{Z}$. By definition, this point is independent of A . Consider a finite subset $\mathcal{Z}' \subset \mathcal{Z}$. The blow-up space $X_{\mathcal{Z}'}$ is homeomorphic to the sphere (for a finite set \mathcal{Z}' , this is both intuitively obvious and technically simple). Let $\sim_{\mathcal{Z}'}$ be the equivalence relation on $X_{\mathcal{Z}'}$, whose non-trivial classes are pullbacks of sets in $\mathcal{Z} - \mathcal{Z}'$ under $\Pi_{\mathcal{Z}'}$. Let $Y_{\mathcal{Z}'}$ be the quotient space by this equivalence relation. By Theorem 15 of Moore, the space $Y_{\mathcal{Z}'}$ is homeomorphic to the sphere. Here, we use our assumption that \mathcal{Z} form a null-sequence (otherwise, the equivalence relation would not be closed). Define the countable subset set $Z_{\mathcal{Z}'} \subset Y_{\mathcal{Z}'}$ as the set of points in $Y_{\mathcal{Z}'}$, whose fibers under the quotient map $X_{\mathcal{Z}'} \rightarrow Y_{\mathcal{Z}'}$ are non-trivial.

There is a canonical projection $\Pi_{\mathcal{Z}, \mathcal{Z}'} : X_{\mathcal{Z}} \rightarrow X_{\mathcal{Z}'}$ mapping an element $\chi : \mathcal{Z} \rightarrow S^2$ to its restriction to \mathcal{Z}' . Denote by $\phi_{\mathcal{Z}'}$ the composition of this projection and the quotient map $X_{\mathcal{Z}'} \rightarrow Y_{\mathcal{Z}'}$.

We can now define elementary curves in $X_{\mathcal{Z}}$. Take a simple curve C in $Y_{\mathcal{Z}'}$ avoiding the set $Z_{\mathcal{Z}'}$. The pullback of C under $\phi_{\mathcal{Z}'}$ is called an elementary curve in $X_{\mathcal{Z}}$. A standard Baire category argument shows that the set of paths in S^2 avoiding a given countable subset of S^2 is dense in the space of all paths with uniform topology. It follows that there are many elementary curves in $X_{\mathcal{Z}}$. Clearly, the set of elementary curves is stable under taking segments and concatenations.

Proposition 13 *The space $X_{\mathcal{Z}}$ satisfies the Elementary domain axiom and the Extension axiom.*

Proof Let \mathcal{Z}' be a finite subset of \mathcal{Z} , and C a simple closed curve in $Y_{\mathcal{Z}'}$ avoiding the set $Z_{\mathcal{Z}'}$. By the Jordan curve theorem, C divides $Y_{\mathcal{Z}'}$ into two connected components, say, U and U' . It follows that the pullbacks of U and U' under $\phi_{\mathcal{Z}'}$ are components of the complement to the elementary curve $\phi_{\mathcal{Z}'}^{-1}(C)$ in $X_{\mathcal{Z}}$. The pullbacks of U and U' are connected because all fibers of $\phi_{\mathcal{Z}'}$ are connected. The Extension axiom for $X_{\mathcal{Z}}$ follows from the corresponding property of $Y_{\mathcal{Z}'}$ (where the elementary curves in $Y_{\mathcal{Z}'}$ are understood as simple curves avoiding the set $Z_{\mathcal{Z}'}$). □

Proposition 14 *Elementary domains form a basis of topology in $X_{\mathcal{Z}}$.*

Proof We need to prove that, for every open set V in $X_{\mathcal{Z}}$ and every point $\chi_0 \in V$, there exists an elementary domain D that contains χ_0 and is contained in V . Moreover, we can assume that V has the following form: there are open sets U_1, \dots, U_n in S^2 and elements $A_1, \dots, A_n \in \mathcal{Z}$ such that V consists of all $\chi \in X_{\mathcal{Z}}$ with the property $\chi(A_i) \in U_i$ for $i = 1, \dots, n$. Recall that such open sets V form a basis in the product topology.

Set $\mathcal{Z}' = \{A_1, \dots, A_n\}$. Clearly, there is an open set V' in $X_{\mathcal{Z}'}$ such that $\chi \in V$ if and only if $\Pi_{\mathcal{Z}, \mathcal{Z}'}(\chi) \in V'$. (The set V' consists of all $\chi' \in X_{\mathcal{Z}'}$ such that $\chi'(A_i) \in U_i$ for $i = 1, \dots, n$.) Set $\chi'_0 = \Pi_{\mathcal{Z}, \mathcal{Z}'}(\chi_0)$. Clearly, there is a Jordan domain neighborhood of χ'_0 contained in V' , whose boundary maps to a simple closed curve in $Y_{\mathcal{Z}'}$ avoiding the set $\mathcal{Z}_{\mathcal{Z}'}$. The corresponding elementary domain is a neighborhood of χ_0 contained in V . \square

Proposition 15 *The space $X_{\mathcal{Z}}$ satisfies the Covering axiom.*

Proof Let Q be an elementary quadrilateral in $X_{\mathcal{Z}}$, and \mathcal{U} an open covering of \overline{Q} . By Proposition 14, it suffices to consider the case, where all elements of \mathcal{U} are elementary domains. By compactness, we can also assume that \mathcal{U} is finite. For every $U \subset \mathcal{U}$, the curve $\Pi_{\mathcal{Z}}(\partial U)$ intersects only finitely many elements of \mathcal{Z} . Let \mathcal{Z}' be a finite subset of \mathcal{Z} such that $\Pi_{\mathcal{Z}}(\partial U)$ are disjoint from $\mathcal{Z} - \mathcal{Z}'$ for all $U \in \mathcal{U}$. The Covering axiom for $X_{\mathcal{Z}}$ now follows from the corresponding statement in $Y_{\mathcal{Z}'}$. \square

Theorem 16 now follows from Theorem 14.

6.3 Existence of regluings

In this section, we prove Theorem 3. Let \mathcal{A} be a countable set of disjoint α -curves in S^2 such that $\alpha(S^1), \alpha \in \mathcal{A}$, form a null-sequence. Consider the set $\mathcal{Z} = \text{Im } \mathcal{A}$. This set satisfies all assumptions of Sect. 6.2. For every α , define Π_{α} as a continuous extension of $\pi_{\alpha} : \overline{\Delta}_{\infty} \rightarrow S^2$ to the whole sphere S^2 such that $\overline{\Delta}$ is mapped to $\alpha(S^1)$. Let $X_{\mathcal{A}}$ be the blow-up space corresponding to the set \mathcal{Z} and the family of projections Π_{α} . By Theorem 16, the space $X_{\mathcal{A}}$ is homeomorphic to the sphere. Clearly, the ungluing space $\Upsilon_{\mathcal{A}}$ embeds into $X_{\mathcal{A}}$ as a closed subset (we will identify $\Upsilon_{\mathcal{A}}$ with its image under this embedding). This proves the first part of Theorem 3. Any component of the complement to $\Upsilon_{\mathcal{A}}$ in $X_{\mathcal{A}}$ is a Jordan domain consisting of all points $\chi \in X_{\mathcal{A}}$ such that $\chi(\alpha) \in \Delta$ for some fixed α . Thus components of $X_{\mathcal{A}} - \Upsilon_{\mathcal{A}}$ are in one-to-one correspondence with the set \mathcal{A} . Let D_{α} denote the component of $X_{\mathcal{A}} - \Upsilon_{\mathcal{A}}$ corresponding to an α -path $\alpha \in \mathcal{A}$.

Recall (Sect. 2.1) that there is a natural identification h_{α} between the boundary of D_{α} ($= S^1_{\alpha}$) and the unit circle S^1 . The composition of $h_{\alpha} : S^1_{\alpha} \rightarrow S^1$ with the projection of S^1 to the t_2 -axis extends to a continuous map

$P_\alpha : \overline{D}_\alpha \rightarrow [-1, 1]$ in such a way that every fiber of this map is a simple curve with endpoints in S^1_α . We can now define an equivalence relation $\sim_{\mathcal{A}}$ on $X_{\mathcal{A}}$ as follows. Two different points of $X_{\mathcal{A}}$ are equivalent, if they lie in the same fiber of some $P_\alpha, \alpha \in \mathcal{A}$. It is easy to see that the equivalence relation $\sim_{\mathcal{A}}$ is closed, and its equivalence classes are connected and non-separating. Therefore, by Theorem 14, the quotient space is homeomorphic to the sphere. In other terms, there is a quotient map $\Psi : X_{\mathcal{A}} \rightarrow S^2$, whose fibers are exactly equivalence classes with respect to $\sim_{\mathcal{A}}$.

Note that, for every $\alpha \in \mathcal{A}$, the map $\beta = \Psi \circ h_\alpha^{-1}$ is a β -path. Let \mathcal{B} be the set of all such β -paths. (We have a fixed one-to-one correspondence between \mathcal{A} and \mathcal{B} .) It is not hard to verify that the curves $\beta(S^1), \beta \in \mathcal{B}$, form a null-sequence. Since the restriction of Ψ to $\Upsilon_{\mathcal{A}}^\circ$ is one-to-one, and, for every $\alpha \in \mathcal{A}$, the map $\Psi \circ h_\alpha^{-1} : S^1 \rightarrow S^2$ equals to the β -path β corresponding to α , we obtain that the map

$$S^2 - \text{Im } \mathcal{A} \xrightarrow{\pi_{\mathcal{A}}^{-1}} \Upsilon_{\mathcal{A}}^\circ \xrightarrow{\Psi} S^2 - \text{Im } \mathcal{B}$$

is a regluing of \mathcal{A} into \mathcal{B} . This concludes the proof of Theorem 3.

6.4 Branched coverings

Let X be a topological sphere and $f : X \rightarrow X$ a continuous map. Recall that f is called a *branched covering* at $x \in X$, if there exist neighborhoods U of x and V of $f(x)$ and homeomorphisms $\phi : U \rightarrow \Delta$ and $\psi : V \rightarrow \Delta$ such that $\psi \circ f \circ \phi^{-1}$ coincides with $z \mapsto z^k$ on Δ , where k is some positive integer. The number k is called the *local degree* of f at x . The map f is a branched covering if it is a branched covering locally at every point. The following is a topological criterion for f to be a branched covering.

Theorem 17 *Suppose that $y = f(x)$, and there exist simply connected domains $U \ni x$ and $V \ni y$ such that $f : U - \{x\} \rightarrow V - \{y\}$ is a covering of degree k . Then f is a branched covering of degree k at x .*

Proof Both $U - \{x\}$ and $V - \{y\}$ can be homeomorphically identified with the quotient of the upper half-plane $\mathbb{H} = \{\text{Im } z > 0\}$ by the translation $z \mapsto z + 1$. Moreover, one can choose the homeomorphisms with the following property: as a point in \mathbb{H}/\mathbb{Z} tends to infinity (i.e. the imaginary part tends to infinity), the corresponding point in $U - \{x\}$ (respectively, in $V - \{y\}$) tends to x (respectively, to y). The map f restricted to $U - \{x\}$ lifts to a homeomorphism $F : \mathbb{H} \rightarrow \mathbb{H}$ such that $F(z + 1) = F(z) + k$. Consider the maps $\Phi(z) = \exp(2\pi i F(z)/k)$ and $\Psi(z) = \exp(2\pi iz)$. These maps descend to

homeomorphisms $\phi : U - \{x\} \rightarrow \Delta - \{0\}$ and $\psi : V - \{y\} \rightarrow \Delta - \{0\}$, respectively. Moreover, we have $\psi \circ f = \phi^k$. Clearly, ϕ and ψ extend continuously to x and y , respectively, and $\phi(x) = \psi(y) = 0$ so that the equality $\psi \circ f = \phi^k$ still holds. □

Corollary 2 *Let $f : S^2 \rightarrow S^2$ be a continuous map and $C \subset S^2$ a finite subset such that $f^{-1}(f(C))$ is finite and f is locally injective on $S^2 - C$. Then f is a branched covering.*

Recall that Theorem 4 states a simple condition that guarantees that a topological regluing of a branched covering is also a branched covering. We are now ready to prove this theorem.

Proof of Theorem 4 Consider a branched covering $f : S^2 \rightarrow S^2$ and a strongly f -stable set \mathcal{A} of disjoint α -paths. Let $\Phi : S^2 - \text{Im } \mathcal{A} \rightarrow S^2 - \text{Im } \mathcal{B}$ be a regluing of \mathcal{A} into a set \mathcal{B} of β -paths. The map $\Phi \circ f \circ \Phi^{-1}$ extends to a continuous map $g : S^2 \rightarrow S^2$, by Theorem 2. We want to prove that g is a branched covering.

Define C_g as the set consisting of points $\Phi(c)$, where $c \notin \text{Im } \mathcal{A}$ is a critical point of f , and of points $\beta(1, 0)$, where $\beta \in \mathcal{B}$ is the β -path corresponding to some α -path $\alpha \in \mathcal{A}$, and $\alpha(0, 1)$ is a critical point of f . Clearly, C_g is finite (its cardinality is the number of critical points of f). By Corollary 2, it suffices to prove that for every $y \notin C_g$ there exists a neighborhood of y , on which g is injective.

First, assume that $y \notin \text{Im } \mathcal{B}$. Then $y = \Phi(x)$ for some point $x \notin \text{Im } \mathcal{A}$. Consider a small neighborhood V of $f(x)$ that does not contain critical values of f . Let U be a Jordan neighborhood of x , whose boundary is disjoint from $\text{Im } \mathcal{A}$ and which is contained in $f^{-1}(V)$. It is not hard to see that g is injective on the Jordan neighborhood of y bounded by $\Phi(\partial U)$.

Next, assume that $y \in \beta(S^1)$, where $\beta \in \mathcal{B}$ is the β -path corresponding to some α -path $\alpha \in \mathcal{A}$, and $\alpha(S^1)$ contains no critical points of f . Consider a small neighborhood V of $f(\alpha(S^1))$ containing no critical values of f . Let U be a Jordan neighborhood of $\alpha(S^1)$, whose boundary is disjoint from $\text{Im } \mathcal{A}$ and which is contained in $f^{-1}(V)$. It is not hard to see that g is injective on the Jordan neighborhood of y bounded by $\Phi(\partial U)$.

Finally, assume that $y = \beta(t_1, t_2)$, where $\beta \in \mathcal{B}$ is the β -path corresponding to some α -path $\alpha \in \mathcal{A}$, and $\alpha(0, 1)$ is a critical point of f . Then we necessarily have $t_2 \neq 0$. Choose a Jordan domain V disjoint from the critical values of f and such that ∂V intersects $f \circ \alpha(S^1)$ exactly by endpoints (so that the critical value $f \circ \alpha(0, 1)$ is in ∂V). Let U_1 and U_2 be the pullbacks of V intersecting $\alpha(S^1)$. These are Jordan domains, whose boundaries intersect exactly by the critical point $\alpha(0, 1)$. By a suitable choice of V , we can arrange that ∂U_1 and ∂U_2 do not intersect $\text{Im } \mathcal{A} - \alpha(S^1)$. The union of $\Phi(\partial U_1 - \{\alpha(0, 1)\})$,

$\Phi(\partial U_2 - \{\alpha(0, 1)\})$ and the endpoints of β is a pair of simple closed curves intersecting exactly by $\beta(1, 0)$. It is easy to see that g is injective on the Jordan neighborhood of y bounded by one of these curves. \square

Acknowledgements I am grateful to Mary Rees for hospitality and very useful conversations during my visit to Liverpool, to K. Pilgrim for a stimulating question that has led to an improvement of results in Sect. 6, to M. Lyubich, J. Milnor, N. Selinger for useful discussions, and to the anonymous referees for the suggested corrections and improvements of the text.

References

1. Aspenberg, M., Yampolsky, M.: Mating non-renormalizable quadratic polynomials. [arXiv:math.DS/0610343](https://arxiv.org/abs/math/0610343)
2. Bing, R.H.: The Kline sphere characterization problem. *Bull. Am. Math. Soc.* **52**, 644–653 (1946)
3. Kuratowski, K.: *Topology*. San Diego, Academic/PWN Press (1966). Revised edition
4. Lei, T., Shishikura, M.: An alternative proof of Mañé's theorem on non-expanding Julia sets. In: *The Mandelbrot Set, Theme and Variations*. LMS Lecture Note Series, vol. 274, pp. 265–279. Cambridge University Press, Cambridge (2000)
5. Luo, J.: *Combinatorics and Holomorphic Dynamics: Captures, Matings and Newton's Method*. PhD Thesis, Cornell University (1995)
6. Mañé, R.: On a Theorem of Fatou. *Bol. Soc. Bras. Mat.* **24**(1), 1–11 (1993)
7. Mañé, R., Sud, P., Sullivan, D.: On the dynamics of rational maps. *Ann. Sci. École Norm. Sup.* (4) **16**(2), 193–217 (1983)
8. Milnor, J.: Geometry and dynamics of quadratic rational maps. *Exp. Math.* **2**, 37–83 (1993)
9. Milnor, J.: *Dynamics in One Complex Variable*, 3rd edn. Princeton University Press, Princeton (2006)
10. Moore, R.L.: On the foundations of plane analysis situs. *Trans. AMS* **17**, 131–164 (1916)
11. Moore, R.L.: Concerning upper-semicontinuous collections of continua. *Trans. AMS* **27**(4), 416–428 (1925)
12. Rees, M.: Components of degree two hyperbolic rational maps. *Invent. Math.* **100**, 357–382 (1990)
13. Rees, M.: A partial description of the parameter space of rational maps of degree two: Part 1. *Acta Math.* **168**, 11–87 (1992)
14. Rees, M.: Views of parameter space, topographer and resident. *Astérisque* **288** (2003)
15. Rees, M.: A fundamental domain for V_3 . Preprint
16. Stimpson, J.: Degree two rational maps with a periodic critical point. PhD Thesis, University of Liverpool, Juli (1993)
17. Timorin, V.: External boundary of M_2 . In: *Fields Institute Communications, Volume 53: Holomorphic Dynamics and Renormalization A Volume in Honour of John Milnor's 75th Birthday*
18. Timorin, V.: On partial semi-conjugacies of quadratic polynomials. Preprint
19. Timorin, V.: Moore's theorem. Preprint
20. van Kampen, E.R.: On some characterizations of 2-dimensional manifolds. *Duke Math. J.* **1**, 74–93 (1935)
21. Zippin, L.: A study of continuous curves and their relation to the Janiszewski–Mullikin theorem. *Trans. Am. Math. Soc.* **31**, 744–770 (1929)