

# “Quasi-Planck” spectra of capillary turbulence on the surface of liquid hydrogen

*M. Yu. Brazhnikov<sup>1</sup>, L. V. Abdurakhimov, S. V. Filatov, A. A. Levchenko*

*Institute of Solid State Physics RAS, 142432 Chernogolovka, Russia*

Submitted 18 November 2010

We report the first observation of “quasi-Planck” spectra of capillary turbulence on the surface of liquid hydrogen in the dissipation domain. Capillary waves have been driven by low-frequency random force. We have observed that the frequency spectrum of surface elevation changes its dependence from power-like  $\langle |\eta_\omega^2| \rangle \sim \omega^{-2.8}$  at middle-frequency domain to “quasi-Planck” distribution  $\sim e^{-\omega/\omega_d}$  at higher frequencies. The frequency  $\omega_d$  is proportional to the boundary frequency between inertial interval and dissipation domain and it is scaled up with the increase of driving force.

*Introduction.* The theory of weak turbulence [1] describes numerous systems of weakly interacting waves: ripples on water and gravity waves on the surface of oceans, Rossby waves in the atmosphere of planets and in the World ocean, Langmuir waves in plasma and spin waves in magnetics. Relatively low viscosity of liquid hydrogen, its low density and the possibility to excite waves on the charged surface by electrical force reveal a unique opportunity for experimental studies of weak wave turbulence. The use of liquid hydrogen for experiments on wave turbulence has already allowed us to study phenomena predicted by the theory, e.g. Kolmogorov-Zakharov steady state spectra of capillary turbulence in a wide range of frequencies [2], as well to observe new ones which have been explained successfully in the framework of the weak turbulence approximation: quasiadiabatic decay of capillary turbulence [3] and suppression of high-frequency turbulent oscillations by additional low-frequency driving force [4].

The weak wave theory predicts an existence of stationary non-equilibrium state in a system of interacting capillary waves – Kolmogorov-Zakharov spectrum for the spectral density of energy  $\varepsilon(k)$  of waves inside inertial interval [5]:

$$\varepsilon(k) \sim P^{1/2} k^{-7/4}. \quad (1)$$

Using relation between the spectral density  $\varepsilon(k)$  and pair correlation function of the surface elevation  $\eta(\mathbf{r}, t)$ , and taking into account the dispersion law for capillary waves  $\omega(k) \sim k^{3/2}$ , one can obtain the frequency spectrum of correlation function  $\langle \eta(t + \tau)\eta(t) \rangle$ :

$$\langle |\eta_\omega^2| \rangle \sim (\sigma k^2)^{-1} \varepsilon(k) (d\omega/dk)^{-1} \sim P^{1/2} \omega^{-17/6}. \quad (2)$$

This stationary spectrum is characterized by a single quantity  $P$  – the energy flux directed towards large fre-

quencies  $\omega$ . At high frequencies the energy transferred is dissipated due to viscous losses and the turbulent cascade is destroyed, therefore to sustain distribution (2) or (1) constant in time the system must be permanently pumped with energy at low frequencies. The high-frequency boundary of the inertial interval  $\omega_d$ , where the distribution (2) is still valid, can be estimated from assumption that the time of viscous damping  $\tau_\nu(\omega)$  and the time of nonlinear interaction  $\tau_{nl}(\omega)$  are of the same order of magnitude for waves at the frequency  $\omega_d$ , this gives us [5]:

$$\omega_d \sim \left( \frac{P^{1/2}}{\nu} \right)^{6/5} \sim \left( \frac{\eta_0^2 \omega_0^{17/6}}{\nu} \right)^{6/5}, \quad (3)$$

where  $\eta_0^2$  is mean squared amplitude of the wave at some low frequency  $\omega_0$ ,  $\nu$  is kinematic viscosity of liquid.

At frequencies above  $\omega_d$  the spectrum depends both on details of dissipation and nonlinear interaction. When waves in dissipation region  $\omega \gg \omega_d$  interact mainly with each other than with waves from inertial range  $\omega \ll \omega_d$  the wave distribution should be close to exponential one [6]. Detailed consideration gives the “quasi-Planck” spectrum for dissipation tail of the correlation function [7]:

$$\langle |\eta_\omega^2| \rangle \sim \omega^{-s} e^{-\omega/\omega_d}, \quad (4)$$

where  $s$  is some constant. The numerical simulations for discrete kinetic equation [7] has confirmed exponential dependence of the wave occupation numbers in the domain of strong damping.

Here we report the first experimental observations of turbulent spectra of the capillary waves in dissipation region driven by low frequency random force on the surface of liquid hydrogen.

*Experimental techniques.* The experimental setup consists of optical cell placed inside the vacuum space

<sup>1</sup>) e-mail: makc@issp.ac.ru

of helium cryostat and optical system for registration of oscillations on the surface of liquids. A conductive cylindrical vessel 6 mm in depth and 60 mm of inner diameter was mounted inside the cell and a flat conductive plate was fixed at 4 mm above the vessel. Gaseous hydrogen has been condensed into the vessel. Radioactive plate located at the bottom of vessel has ionised the liquid hydrogen. In the presence of dc voltage of about 1 kV between the vessel and upper plate ions have been separated and charges with the sign corresponding to the polarity of dc voltage have been accumulated under the surface of liquid hydrogen. Additional ac voltage up to maximum value 100 V was used to excite waves on the surface of liquid.

The use of electric field for driving capillary waves makes a great advantage comparing with the usual techniques for exciting waves on the water surface, e.g. via Faraday instability. It allows us to drive the surface accurately by sufficiently well controllable force. In this set of experiments we have used low-frequency random signals as a driving ac voltage. The signals have been synthesised via Fourier transform from random set of phases and a boxcar like amplitude spectrum, which is non-zero only inside some frequency interval (pumping range).

We use the laser reflection method to detect waves on the surface of liquid. A laser beam incident at small sliding angle (about 0.2 rad) on the surface of liquid is reflected and focused by lens onto photo-detector. The voltage from photo-detector is amplified and digitised by 24-bit analog to digital converter with sampling frequency of about 100 kHz. Waves are detected in regime of “wide beam”, when size of the laser spot is greater than typical wave length. The power of reflected light  $P(t)$  in this regime is proportional to the surface elevation  $\eta(t)$  [8]. For this reason we will not make the difference further for spectra of the correlation function of surface elevation  $\langle |\eta_\omega^2| \rangle$  and the reflected beam power  $\langle |P_\omega^2| \rangle$ .

The maximum wave steepness that can be registered in our experimental setup is limited by sizes of optical windows in cryostat and it is approximately equal to 0.05.

*Experimental results and discussion.* Capillary waves have been excited by random force in a frequency range 39–103 Hz. The root mean square of driving voltage was varied from  $V_p = 0$  V, i.e. no pumping at all, up to maximum value  $V_p = 30$  V limited mainly by the maximum allowed wave steepness. Fig.1 demonstrates Fourier spectra of reflected laser beam power  $P_\omega^2$  for different amplitudes of driving force. The excitation range can be easily seen at low frequency part of all spectra.




Fig.1. Spectra of surface oscillation  $P_\omega^2$  excited by random force in a frequency range 39–103 Hz at different amplitudes of driving force. The rms of driving voltage  $V_p$  is changed from 4 to 30 V. Darker line corresponds to stronger driving force. The arrow indicates the high-frequency boundary of the inertial interval  $\omega_b \approx 4$  kHz for driving voltage  $V_p = 30$  V

The excitation range is followed by inertial interval – relatively wide frequency domain where spectrum  $P_\omega^2$  exhibits power-law dependence (2). The width of inertial interval depends on drive amplitude. When the surface is excited by weak force  $V_p = 4$  V the dissipation starts nearly from the excitation range and no inertial interval is observed. Increasing the driving force leads to broadening the inertial interval, the high-frequency boundary  $\omega_b$  of inertial interval shifts towards higher frequencies. The widest inertial range from  $\approx 0.3$  kHz to  $\omega_b \approx 4$  kHz is seen for maximum driving voltage  $V_p = 30$  V. At frequencies above the high-frequency boundary surface oscillations degrade due to viscous losses, curves  $P_\omega^2$  go down smoothly and hide finally below the level of apparatus noise.

Turbulent spectra replotted in linear scale in Fig.2 demonstrate that the decrease of wave amplitudes at frequencies above the high-frequency boundary of inertial range can be approximated good enough by the exponential decay  $P_\omega^2 \sim e^{-\omega/\omega_d}$  in some interval. Fitting procedure is consistent with initial assumption  $\omega \gg \omega_d$ , the resulting fitting parameter  $\omega_d$  is considerably less than frequencies from fitting interval. For examples, spectrum at  $V_p = 26$  V is approximated in the range of 5–9 kHz with  $\omega_d \approx 0.6$  kHz. Unfortunately, narrow fitting interval does not allow to establish exponent  $s$  of the “quasi-Planck” (4) distribution accurately. Fitted  $\omega_d$  is also several times less than visible boundary between inertial and dissipation domains (see Fig.2). This discrepancy can be attributed to a certain degree to a freedom




Fig.2. Spectra  $P_\omega^2$  for driving voltage  $V_p = 8$  V (light grey), 16 V (grey) and 26 V (dark grey) in linear scale. Dotted line – power-law distribution  $\sim \omega^{-2.8}$ , dashed lines – fitting functions  $\sim e^{-\omega/\omega_d}$ .  $\omega_d$  is approximately equal to 0.2, 0.4 and 0.6 kHz for  $V_p = 8, 16$  and 26 V correspondingly

in definition of the boundary frequency (3), which can be renormalized by some constant.

The boundary of viscous damping  $\omega_d$  obtained from fitting spectra by exponential decay in dissipation interval grows with the increase of driving force. We use surface response  $\eta_0$ , namely the absolute value of  $P_\omega$  at frequency of 53 Hz (location of the maximum of  $P_\omega^2$  distribution inside excitation range), as a measure for excitation level. The quantity  $\eta_0$  is directly proportional to the mean wave height at the same frequency. Fig.3 shows




Fig.3. Dependence of viscous boundary  $\omega_d$  (black dots) on mean height of low-frequency wave  $\eta_0$ , grey line – fitting function  $\eta_0^{0.85}$

that the boundary frequency can be fitted by power-law function  $\omega_d(\eta_0) \sim \eta_0^m$  with the exponent  $m = 0.85 \pm 0.05$ . It should be noted that fitting the experimental spectra by the “quasi-Planck” with small non-zero  $s$  ( $|s| \leq 2$ )

affects slightly  $\omega_d$  (less than 20%). However this correction does not modify within the fitting error the exponent  $m$  in dependence of the boundary frequency on excitation level.

The observed exponent  $m \approx 0.85$  differs significantly with expected index  $m = 12/5$  from formula (3). It worth to mention that in case of turbulent cascades excited by harmonic force the measured boundary frequency was in good agreement with estimations [9],  $\omega_d(\eta) \sim \eta^{1.3}$ .

**Conclusion.** The transition from power-like Kolmogorov-Zakharov spectrum (2) inside inertial interval to “quasi-Planck” distribution  $\omega^{-s} e^{-\omega/\omega_d}$  in dissipation region for capillary turbulence has been observed for the first time. The exponential decay in dissipation region  $\omega/\omega_d \gg 1$  supports theoretical consideration and are in qualitative agreement with numerical simulations [7]. The boundary of viscous damping  $\omega_d$  grows with increasing the drive amplitude and depends on mean wave height  $\eta_0$  at driving frequency as  $\omega_d \sim \eta_0^{0.85 \pm 0.05}$ . However the observed dependence diverges from estimations, the exponent is almost three times smaller than predicted index.

The work has been supported by the Russian Foundation for Basic Research under grant #09-02-01146-a, by Russian Academy of Sciences (program “Quantum Physics of Condensed Matter”) and by Ministry of education and science of the Russian Federation (grant NSh-4415.2010.2). M.Yu. Brazhnikov acknowledges support from the Russian Science Support Foundation.

1. V. E. Zakharov, V. S. L'vov, and G. Falkovich, *Kolmogorov Spectra of Turbulence I*, Springer-Verlag, 1992.
2. M. Yu. Brazhnikov, G. V. Kolmakov, A. A. Levchenko, and L. P. Mezhev-Deglin, *JETP Lett.* **73**, 398 (2001).
3. G. V. Kolmakov, A. A. Levchenko, M. Yu. Brazhnikov et al., *Phys. Rev. Lett.* **93**, 074501 (2004).
4. M. Yu. Brazhnikov, G. V. Kolmakov, A. A. Levchenko, and L. P. Mezhev-Deglin, *JETP Lett.* **82**, 642 (2005).
5. V. E. Zakharov and N. N. Filonenko, *J. App. Mech. Tech. Phys.* **8**, 62 (1967).
6. V. M. Malkin, *JETP* **86**, 1263 (1984).
7. I. V. Ryzhenkova and G. E. Falkovich, *JETP* **98**, 1931 (1990).
8. M. Yu. Brazhnikov, A. A. Levchenko, and L. P. Mezhev-Deglin, *Instr. Exp. Tech.* **45**, 758 (2002).
9. M. Yu. Brazhnikov, G. V. Kolmakov, A. A. Levchenko, and L. P. Mezhev-Deglin, *JETP Lett.* **74**, 583 (2001).