



ELSEVIER

Contents lists available at ScienceDirect

Journal of Functional Analysis

www.elsevier.com/locate/jfa



Corrigendum

A corrected quantitative version of the Morse lemma



Sébastien Gouëzel^{a,*}, Vladimir Shchur^{b,c}

^a *Laboratoire Jean Leray, CNRS UMR 6629, Université de Nantes, 2 rue de la Houssinière, 44322 Nantes, France*

^b *Departments of Integrative Biology and Statistics, University of California Berkeley, 4098 Valley Life Sciences Building (VLSB), Berkeley, CA 94720-3140, USA*

^c *National Research University Higher School of Economics, 101000 Moscow, Russia*

ARTICLE INFO

Article history:

Received 17 October 2018

Accepted 26 February 2019

Available online 21 March 2019

Communicated by Stefaan Vaes

Keywords:

Morse lemma

Gromov-hyperbolic space

Quasi-geodesic

Isabelle/HOL

ABSTRACT

There is a gap in the proof of the main theorem in the article [5] on optimal bounds for the Morse lemma in Gromov-hyperbolic spaces. We correct this gap, showing that the main theorem of [5] is true. We also describe a computer certification of this result.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

The Morse lemma is a fundamental result in the theory of Gromov-hyperbolic spaces. It asserts that, in a δ -hyperbolic space, the Hausdorff distance between a (λ, C) -quasi-geodesic and a geodesic segment sharing the same endpoints is bounded by a constant $A(\lambda, C, \delta)$ depending only on λ , C and δ , and not on the length of the geodesic. Many

DOI of original article: <https://doi.org/10.1016/j.jfa.2012.11.014>.

* Corresponding author.

E-mail addresses: sebastien.gouezel@univ-nantes.fr (S. Gouëzel), vlshchur@gmail.com (V. Shchur).

<https://doi.org/10.1016/j.jfa.2019.02.021>

0022-1236/© 2019 Elsevier Inc. All rights reserved.

proofs of this result have been given, with different expressions for A . An optimal value for A (up to a multiplicative constant) has only been found recently in the article [5] by the second author, giving $A(\lambda, C, \delta) = K\lambda^2(C + \delta)$ for an explicit constant $K = 4(78 + 133/\log(2) \cdot \exp(157 \log(2)/28)) \sim 37723$.

Unfortunately, there is a gap in the proof of this theorem in [5], which was noticed by the first author while he was developing a library [4] on Gromov-hyperbolic spaces in the computer assistant Isabelle/HOL. In such a process, all proofs are formalized on a computer, and checked starting from the most basic axioms. The degree of confidence reached after such a formal proof is orders of magnitude higher than what can be obtained by even the most diligent reader or referee, and indeed this process shed the light on the gap in [5]. The gap is on Page 829: the inequality $\sum_{i=1}^n e^{-X_i}(X_{i-1} - X_i) \leq \int_0^\infty e^{-t} dt$ goes in the wrong direction as the sequence X_i is decreasing.

In this paper, we fix this gap. Here is the estimate we get.

Theorem 1.1. *Consider a (λ, C) -quasi-geodesic Q in a δ -hyperbolic space X , and G a geodesic segment between its endpoints. Then the Hausdorff distance $HD(Q, G)$ between Q and G satisfies*

$$HD(Q, G) \leq 92\lambda^2(C + \delta).$$

Let us specify precisely the terms used in this statement, as there are small variations in the definitions in the literature. For us, a (λ, C) -quasi-geodesic is the image of a map f from a compact interval to X satisfying for all x, y the inequalities

$$\lambda^{-1} |y - x| - C \leq d(f(x), f(y)) \leq \lambda |y - x| + C.$$

A map satisfying these inequalities is also called a (λ, C) -quasi-isometry. We also require $\lambda \geq 1$ and $C \geq 0$ in the definition. A geodesic segment is by definition a $(1, 0)$ -quasi-geodesic. We say that the space X is δ -hyperbolic if the Gromov product $(x, y)_w = (d(x, w) + d(y, w) - d(x, y))/2$ satisfies for all points x, y, z, w the inequality

$$(x, z)_w \geq \min((x, y)_w, (y, z)_w) - \delta.$$

Finally, the Hausdorff distance $HD(Q, G)$ is the smallest number r such that G is included in the r -neighborhood of Q , and conversely.

Remark 1.2. For any $\lambda \geq 3$, $C \geq 0$ and $\delta \geq 0$, one can construct an example of a (λ, C) -quasi-geodesic Q in a δ -hyperbolic space which satisfies $HD(Q, G) \geq \lambda^2(C + \delta)/9$ where G is a geodesic segment joining the endpoints of Q . This shows that Theorem 1.1 is optimal, up to the value of the multiplicative constant. Such examples for $\delta = 0$ are already given in [5], and the following is a variation around these examples.

Example 1.3. Let $\lambda \geq 3$, $C \geq 0$ and $\delta \geq 0$. Take $X = \mathbb{R} \times [0, \delta]$ with the L^1 distance. This is a δ -hyperbolic space. Let $\bar{\lambda} = \lambda/3 \geq 1$. Define a quasi-geodesic $f : [0, 2\bar{\lambda}(C + \delta) + \delta/\bar{\lambda}] \rightarrow$

X by going always at speed $\bar{\lambda}$ from $(0, 0)$ to $(\bar{\lambda}^2(C + \delta), 0)$, then to $(\bar{\lambda}^2(C + \delta), \delta)$, then to $(0, \delta)$. The Hausdorff distance between the quasi-geodesic Q defined by f and the geodesic G joining $(0, 0)$ and $(0, \delta)$ is $\bar{\lambda}^2(C + \delta) = \lambda^2(C + \delta)/9$. We claim that f is a (λ, C) -quasi-geodesic. The upper bound $d(f(x), f(y)) \leq \lambda|y - x| + C$ is obvious as f is $\bar{\lambda}$ -Lipschitz by construction. For the lower bound $d(f(x), f(y)) \geq \lambda^{-1}|y - x| - C$, the most demanding points are the endpoints of the interval $x = 0$ and $y = 2\bar{\lambda}(C + \delta) + \delta/\bar{\lambda}$: we should check that

$$d(f(x), f(y)) = \delta \geq \lambda^{-1} \cdot (2(C + \delta)\bar{\lambda} + \delta/\bar{\lambda}) - C.$$

This follows from the choice $\bar{\lambda} = \lambda/3$.

The new proof of Theorem 1.1 has been completely formalized in Isabelle/HOL in [4]. Therefore, the above theorem is certified. Here is this statement as proved in Isabelle/HOL.

```
theorem (in Gromov_hyperbolic_space) Morse_Gromov_theorem':
  fixes f: "real  $\Rightarrow$  'a"
  assumes "lambda C-quasi_isometry_on {a..b} f"
    "geodesic_segment_between G (f a) (f b)"
  shows "hausdorff_distance (f`{a..b}) G  $\leq$  92 * lambda^2 * (C + deltaG(TYPE('a)))"
```

In this formal statement, 'a is a type of class `Gromov_hyperbolic_space`. It corresponds to the space X of Theorem 1.1, and the associated hyperbolicity constant is `deltaG(TYPE('a))`. Instead of talking of the quasi-geodesic Q , the formal statement is made in terms of its parametrization f , as the notion of endpoint of a quasi-geodesic is not really well defined. With this correspondence, the two statements directly correspond to each other.

Although the proof is more involved than the original argument in [5], the constant we get in the end is much better (92 instead of 37724). Indeed, we have tried to optimize the constant as much as we could, contrary to [5], keeping in mind the foundational nature of the library [4]. This optimization owes a lot to the formalization process. It makes it possible to optimize locally one part of the proof, and see if it breaks other parts of the proof by checking if the proof assistant complains that the proof is not correct any more, or if everything goes through. The certainty of the result also makes the optimization worth it, as we are sure not to have forgotten for example an edge case that would spoil the estimates.

Having a formalized certified proof raises interesting questions about the way to write mathematics. We do not need to convince a reader (or a referee!) that the result is correct, as we have already done the much more demanding task of convincing a computer, and the proof with all details can be read by the interested reader in [4]. Rather, we have to convey the interesting ideas. We have decided to give all the precise statements we use (in their traditional version, but the very same statements have been formalized

in [4]), but skip their proofs if they are small variations around results that are already available in the literature. For the main proof, we will explain (with as many details as in a traditional mathematical paper) a simplified version of the proof that gives the same statement as Theorem 1.1 but not caring much about the universal constants (this simplified argument gives the constant 2460 instead of 92 in Theorem 1.1). Then we will comment without entering in too many details on the various optimizations that can be done, leading to the above statement.

Remark 1.4. The proof of Theorem 1.1 is delicate. However, we would like to emphasize that this is not due to our desire to formalize the proof on computer: the argument we give in this article is the simplest one we have been able to come up with, without any attempt to get an easy to formalize proof. And indeed this proof was not easy to formalize, but the mere fact that this was possible shows how powerful proof assistants already are today.

2. Proof of the main theorem

The proof uses the notion of quasiconvexity. We say that a subset $Y \subseteq X$ is K -quasiconvex if, for any $y_1, y_2 \in Y$, there exists a geodesic between y_1 and y_2 which is included in the K -neighborhood of Y . For instance, geodesics are 0-quasiconvex. The r -neighborhood of a 0-quasiconvex set is always 8δ -quasiconvex, see [3, Proposition 10.1.2].

We follow the global strategy of [5] to prove Theorem 1.1, with a new more involved argument at a key technical step. Thanks to [1], we can assume without loss of generality that the space X is geodesic. The quasi-geodesic Q is by definition the image of a (λ, C) -quasi-isometric map $f : [u^-, u^+] \rightarrow X$. The statement for a general quasi-isometric map f reduces to the one for a continuous quasi-isometric map f thanks to the following approximation lemma, which is a version of [5, Lemma 9] or [2, Lemma III.H.1.11].

Lemma 2.1. *Consider a (λ, C) -quasi-isometry from a compact interval to a geodesic metric space, whose endpoints are at distance at least $2C$. Then it is within Hausdorff distance $2C$ of a $(\lambda, 4C)$ -quasi-geodesic with the same endpoints which is moreover 2λ -Lipschitz.*

The proof of this lemma is very classical: assume that the initial quasi-geodesic is defined on an interval $[u^-, u^+]$. Then the assumptions ensure that $u^+ - u^- \geq C/\lambda$. Split suitably the interval $[u^-, u^+]$ into subintervals with length in $[C/\lambda, 2C/\lambda]$. The new quasi-geodesic will coincide with the initial one on the endpoints of these subintervals, and be geodesic in between. The facts that this new function is a $(\lambda, 4C)$ -quasi-geodesic, within Hausdorff distance $2C$ of the original one, and 2λ -Lipschitz, follow from direct computations.

Replacing the original quasi-geodesic by the new one given by Lemma 2.1 and C by $4C$, we will assume from this point on that the (λ, C) -quasi-geodesic f is also continuous. Replacing the original hyperbolicity constant δ_0 by a slightly larger constant δ (and letting δ tend to δ_0 at the end of the argument), we can assume that the space is hyperbolic for a constant strictly smaller than δ , and also that $\delta > 0$.

Consider $z \in [u^-, u^+]$. We want to estimate $d(f(z), G)$. We will prove an estimate of the form

$$d(f(z), G) \leq K_0 + \frac{K_1}{K_2} \int_0^{u^+ - u^-} e^{-K_2 t} dt = K_0 + K_1 \cdot (1 - e^{-K_2(u^+ - u^-)}), \tag{2.1}$$

where K_0, K_1 and K_2 are suitable parameters that do not depend on u^- and u^+ . Both K_0 and K_1 will be of the form $K_i = k_i \lambda^2(C + \delta)$, while K_2 will be of the form $K_2 = k_2 / (\delta \lambda)$ where k_0, k_1, k_2 are explicit positive real constants. They will be defined in (2.4), (2.7) and (2.6). This estimate is proved inductively over the size of $u^+ - u^-$, reducing the estimate over $[u^-, u^+]$ to the estimate over a shorter interval $[v^-, v^+]$. We will have to show that the loss in this reduction process is controlled in terms of $K_1 e^{-K_2(v^+ - v^-)} - K_1 e^{-K_2(u^+ - u^-)}$, to conclude the proof of (2.1) by induction.

Let us first explain why this estimate concludes the proof. It implies that $d(f(z), G) \leq K_0 + K_1$. This proves that the image Q of f is included in the $(k_0 + k_1)\lambda^2(C + \delta)$ -neighborhood of G . To get the estimate on the Hausdorff distance, one needs to show that G is also included in a $k\lambda^2(C + \delta)$ -neighborhood of Q for some k . This follows from the previous estimate and a standard argument (see [2]) that we recall now. Consider a point $g \in G$. Denote by Q^- the set of points on Q that are within distance $(k_0 + k_1)\lambda^2(C + \delta)$ of a point of G in $[f(u^-), g]$, and by Q^+ the set of points on Q that are within distance $(k_0 + k_1)\lambda^2(C + \delta)$ of a point of G in $[g, f(u^+)]$. The previous estimate implies that $Q = Q_1 \cup Q_2$. As Q is connected, it follows that $Q_1 \cap Q_2 \neq \emptyset$. Denote by $f(z)$ a point in this intersection, and by g^- and g^+ two points before and after g on G , at distance at most $(k_0 + k_1)\lambda^2(C + \delta)$ of $f(z)$. Using hyperbolicity in a triangle with vertices at $g^-, g^+, f(z)$ and the fact that g is on a geodesic between g^- and g^+ , it follows that the distance between g and $f(z)$ is at most $(k_0 + k_1)\lambda^2(C + \delta) + \delta$. As $\lambda \geq 1$, this expression is bounded by $(k_0 + k_1 + 1)\lambda^2(C + \delta)$. This concludes the argument, for the constant $k = k_0 + k_1 + 1$. We remind that [6] contains a stronger result (Theorem 3) claiming that the geodesic G is included in an $A(\delta \log \lambda + C + \delta)$ -neighborhood of the quasi-geodesic Q with some universal constant A .

It remains to prove the estimate (2.1). The proof will use two parameters L and D . For simplicity, let us take

$$L = D = 100\delta. \tag{2.2}$$

We keep separate notations for L and D because we will want to optimize the choice of their values later.

Case 1. The case where $d(f(z), G) \leq L$ is trivial, as the estimate (2.1) holds if one takes K_0 large enough.

Case 2. Let us therefore assume $d(f(z), G) > L$. We will construct several points along $[u^-, z]$. To ease the reading, their order will correspond to the alphabetical order when possible.

Consider a projection π_z of $f(z)$ on G , and a geodesic segment H from π_z to $f(z)$. Denote by $p : X \rightarrow H$ a closest-point projection on H . The idea is to project the quasi-geodesic Q on H and to consider the subpart Q' of Q that projects at distance at least L of π_z . If one could show that Q' is quantitatively shorter than Q and that the distance from $f(z)$ to π_z is controlled in terms of the distance from $f(z)$ to a geodesic joining the endpoints of Q' , then we would be in good shape to prove (2.1) inductively, deducing the estimate for Q from the estimate for Q' . The real argument will be built around this naive idea, but in a more subtle way.

More precisely, consider two points $y^- \in [u^-, z]$ and $y^+ \in [z, u^+]$ such that the projections $p(f(y^-))$ and $p(f(y^+))$ are at distance roughly L of π_z . In general, p is not uniquely defined and not continuous, but this is almost the case up to $O(\delta)$ thanks to the hyperbolicity of the space. With the following standard lemma and recalling that H is 0-quasiconvex as it is a geodesic, one can find y^- and y^+ such that

$$d(p(f(y^\pm)), \pi_z) \in [L - 4\delta, L]. \tag{2.3}$$

Lemma 2.2. *A closest-point projection of a connected set on a K -quasiconvex subset Y of X has gaps of size at most $4\delta + 2K$. More precisely, if $f : [a, b] \rightarrow X$ is a continuous function and $p(f(t))$ denotes a closest point projection of $f(t)$ on Y , then for any $\tau \leq d(p(f(a)), p(f(b)))$, there exists $t \in [a, b]$ such that $d(p(f(a)), p(f(t))) \in [\tau - 4\delta - 2K, \tau]$. Moreover, one can ensure that $d(p(f(a)), p(f(s))) \leq d(p(f(a)), p(f(t)))$ for all $s \leq t$.*

Denote by d^- (respectively d^+) the minimal distance of a point in $f([u^-, y^-])$ (respectively $f([y^+, u^+])$) to H . These distances are realized by two points $f(m^-)$ and $f(m^+)$, by continuity of f .

Case 2.1. Assume that $\max(d^-, d^+)$ is not large, say $\leq D + C$ where $D = 100\delta$ is the constant we have chosen in (2.2) and C is the quasi-isometry parameter. This is again an easy case. Indeed, as the projections of $f(m^-)$ and $f(m^+)$ are within distance L of π_z , one gets $d(f(m^-), f(m^+)) \leq 2D + 2C + L$. By quasi-isometry,

$$d(m^-, m^+) \leq \lambda(d(f(m^-), f(m^+)) + C) \leq \lambda(2D + 3C + L).$$

As z is between m^- and m^+ , one gets in particular $d(m^-, z) \leq \lambda(2D + 3C + L)$. Then

$$\begin{aligned} d(f(z), \pi_z) &\leq d(f(z), f(m^-)) + d(f(m^-), p(f(m^-))) + d(p(f(m^-)), \pi_z) \\ &\leq (\lambda d(z, m^-) + C) + (D + C) + L \leq \lambda^2(3D + 5C + 2L). \end{aligned}$$

This is compatible with the inequality (2.1) if one takes

$$K_0 = 500\lambda^2(\delta + C). \tag{2.4}$$

Case 2.2. Assume now that $\max(d^-, d^+) \geq D + C$, and $d^- \geq d^+$ for instance. This is the interesting case. The main step in the proof is the following lemma.

Lemma 2.3. *There exist two points $v \leq x$ in $[u^-, y^-]$ and a real number $d' \geq d^-$ such that*

$$L - 74\delta \leq 4\sqrt{2}\lambda(x - v)e^{-d' \log(2)/(10\delta)} \tag{2.5}$$

and $d(f(v), p(f(v))) \leq 4d'$.

The numerology in the lemma (74 and $4\sqrt{2}$ and $\log(2)/10$ and 4) is of no importance: what only matters is that $L - 74\delta$ is positive, thanks to the choice of L in (2.2), and that the other numbers are positive and fixed.

Let us show how to conclude the proof using the lemma. We have

$$\begin{aligned} m^+ - v &= d(v, m^+) \leq \lambda(d(f(v), f(m^+)) + C) \\ &\leq \lambda\left(d(f(v), p(f(v))) + d(p(f(v)), p(f(m^+))) + d(p(f(m^+)), f(m^+)) + C\right) \\ &\leq \lambda(4d' + L + d^+ + C) \leq 6\lambda d', \end{aligned}$$

as $L + C = D + C \leq d^- \leq d'$ and $d^+ \leq d^- \leq d'$. Therefore, taking

$$K_2 = \log(2)/(60\delta\lambda), \tag{2.6}$$

the inequality (2.5) gives

$$\begin{aligned} L - 74\delta &\leq 4\sqrt{2}\lambda(x - v)e^{-(m^+ - v) \cdot \log(2)/(60\delta\lambda)} = \frac{4\sqrt{2}\lambda}{K_2} \cdot K_2(x - v)e^{-K_2(m^+ - v)} \\ &\leq \frac{4\sqrt{2}\lambda}{K_2}(e^{K_2(x - v)} - 1)e^{-K_2(m^+ - v)} = \frac{4\sqrt{2}\lambda}{K_2}(e^{-K_2(m^+ - x)} - e^{-K_2(m^+ - v)}) \\ &\leq \frac{4\sqrt{2}\lambda}{K_2}(e^{-K_2(m^+ - x)} - e^{-K_2(u^+ - u^-)}). \end{aligned}$$

Consider a new geodesic G' between $f(x)$ and $f(m^+)$. Arguing by induction, we can assume that the estimate (2.1) has already been proved for G' , and we want to deduce it for G . Since both endpoints of G' project within distance L of π_z , one checks that the distance from $f(z)$ to G is controlled by the distance from $f(z)$ to G' (this is a version of [5, Lemma 5]). More specifically,

$$d(f(z), G) \leq d(f(z), G') + L + 4\delta.$$

Bounding $d(f(z), G')$ thanks to the induction assumption, and plugging in the estimate from the previous equation, we get

$$d(f(z), G) \leq K_0 + K_1(1 - e^{-K_2(m^+ - x)}) + \frac{L + 4\delta}{L - 74\delta} \cdot \frac{4\sqrt{2}\lambda}{K_2}(e^{-K_2(m^+ - x)} - e^{-K_2(u^+ - u^-)}).$$

Let us take

$$K_1 = \frac{L + 4\delta}{L - 74\delta} \cdot \frac{4\sqrt{2}\lambda}{K_2}. \tag{2.7}$$

Then the terms $K_1 e^{-K_2(m^+ - x)}$ simplify in this equation, and we are left with

$$d(f(z), G) \leq K_0 + K_1(1 - e^{-K_2(u^+ - u^-)}).$$

This is (2.1), as desired. This concludes the proof of Theorem 1.1. \square

It remains to prove Lemma 2.3. The argument relies on the contracting properties of closest-point projections on quasiconvex sets. The first such basic statement is the following variation around [3, Proposition 10.2.1].

Lemma 2.4. *Consider a K -quasiconvex subset Y of X . Then projections p_x and p_y on Y of two points x and y satisfy*

$$d(p_x, p_y) \leq \max(5\delta + 2K, d(x, y) - d(x, p_x) - d(y, p_y) + 10\delta + 4K).$$

This result expresses the classical fact that a geodesic from x to y essentially follows a geodesic from x to p_x , then from p_x to p_y , then from p_y to y .

The second result we need is more sophisticated. Instead of a linear gain in terms of the distance to the set one projects on, as in the previous lemma, it gives an exponential gain in the upper bound, by a successive reduction process. It is proved by putting points along the path with gaps of size 10δ . Then, move by 5δ towards Y : this reduces the distance between the points by 5δ essentially thanks to the previous lemma. Then, discard half the points: this shows that by moving towards Y by 5δ the length of the path has been divided by 2. One can iterate this argument to get the exponential gain. We give a statement for the projection on quasiconvex sets as this is what we will need later on. This statement is proved in [5, Lemma 10] for the projection on a geodesic segment, but the case of a general quasiconvex set is analogous.

Lemma 2.5. Consider a (λ, C) -quasi-geodesic path $f : [a, b] \rightarrow X$, everywhere at distance at least D of a K -quasiconvex subset Y . Then, if $D \geq 15/2 \cdot \delta + K + C/2$, projections p_a of $f(a)$ and p_b of $f(b)$ on Y satisfy the inequality

$$d(p_a, p_b) \leq 2K + 8\delta + \max \left(5\delta, 4\sqrt{2}\lambda(b - a) \exp \left(-(D - K - C/2) \log(2)/(5\delta) \right) \right).$$

Using these results, we can prove Lemma 2.3.

Proof of Lemma 2.3. For $k \geq 0$, let V_k denote the $(2^k - 1)d^-$ -neighborhood of H . These sets are all 8δ -quasiconvex. We recall that $p(f(x))$ is a projection of $f(x)$ on H . Let $p_k(x)$ denote the point on a fixed geodesic between $p(f(x))$ and $f(x)$ at distance $\min((2^k - 1)d^-, d(p(f(x)), f(x)))$ of $p(f(x))$. Then $p_k(x)$ is a projection of $f(x)$ on V_k , and moreover these projections are compatible in the following sense: for $k \leq \ell$, then $p_k(x)$ is a projection of $p_\ell(x)$ on V_k . Moreover, $p_0(x) = p(f(x))$.

We will do an inductive construction over k . This construction will have to stop at some step, where it will give the desired points. Until the argument stops, we will construct a point $x_k \in [u^-, y^-]$ such that

$$d(p_k(u^-), p_k(x_k)) \geq L - 8\delta \tag{2.8}$$

and

$$\text{for all } w \in [u^-, x_k], d(f(w), p_0(w)) \geq (2^{k+1} - 1)d^-. \tag{2.9}$$

Let us first check that this property holds for $k = 0$. Take $x_0 = y^-$. The point π_z is a projection of $f(z)$ on the geodesic G between $f(u^-)$ and $f(u^+)$. This does not imply that the projection $p_0(u^-)$ of $f(u^-)$ on the geodesic H between π_z and $f(z)$ is exactly at π_z (contrary to the situation in the Euclidean plane), but by hyperbolicity one checks that $d(\pi_z, p_0(u^-)) \leq 4\delta$ (this is a version of [5, Lemma 3]). Since $d(\pi_z, p_0(y^-)) \in [L - 4\delta, L]$ by (2.3) and $x_0 = y^-$, we deduce that $d(p_0(u^-), p_0(x_0)) \geq L - 8\delta$. This is (2.8). Moreover, by definition of d^- , the inequality (2.9) holds for $k = 0$.

Assume now that (2.8) and (2.9) hold at k . We will show that either we can find a pair of points that satisfy the conclusion of the lemma, or we can construct a point x_{k+1} such that (2.8) and (2.9) hold at $k + 1$.

As V_k is 8δ -quasiconvex, we deduce from Lemma 2.2 that the gaps of the closest-point projection p_k are bounded by 20δ . Therefore, we can find a point $x_{k+1} \in [u^-, x_k]$ whose projection on V_k satisfies

$$d(p_k(u^-), p_k(x_{k+1})) \in [22\delta, 42\delta], \tag{2.10}$$

and moreover all points $w \in [u^-, x_{k+1}]$ satisfy

$$d(p_k(u^-), p_k(w)) \leq 42\delta. \tag{2.11}$$

There are two cases to consider:

If there exists $v \in [u^-, x_{k+1}]$ with $d(f(v), p_0(v)) \leq (2^{k+2} - 1)d^-$. Then we claim that the pair (v, x_k) satisfies the conclusion of Lemma 2.3, for $d' = 2^k d^-$. First, the inequalities $d' \geq d^-$ and $d(f(v), p_0(v)) \leq 4d'$ hold by construction. Moreover, $d(p_k(v), p_k(x_k)) \geq L - 50\delta$ as $p_k(x_k)$ is far from $p_k(u^-)$ by (2.8), and $p_k(v)$ is close to $p_k(u^-)$ by (2.11). As all intermediate points are at distance at least $(2^{k+1} - 1)d^-$ of V_0 by (2.9), they are at distance at least $2^k d^-$ of V_k and we can apply the exponential contraction Lemma 2.5 with $D = 2^k d^-$. As V_k is 8δ -quasiconvex, we get

$$L - 50\delta \leq d(p_k(v), p_k(x_k)) \leq 24\delta + \max\left(5\delta, 4\sqrt{2}\lambda(x_k - v) \exp\left(- (2^k d^- - 8\delta - C/2) \log(2)/(5\delta)\right)\right).$$

As $L - 50\delta > 29\delta$, the maximum has to be realized by the second term. Moreover, $2^k d^- - 8\delta - C/2 \geq (2^k d^-)/2 = d'/2$, as $d^- \geq D + C = 100\delta + C$. We obtain

$$L - 74\delta \leq 4\sqrt{2}\lambda(x_k - v) \exp\left(-d' \log(2)/(10\delta)\right). \tag{2.12}$$

This concludes the proof in this case.

Otherwise, $d(f(w), p_0(w)) \geq (2^{k+2} - 1)d^-$ for all $w \in [u^-, x_{k+1}]$. In this case, (2.9) holds for $k + 1$. Let us check that (2.8) also holds for $k + 1$, by applying the projection Lemma 2.4 to the points $p_{k+1}(u^-)$ and $p_{k+1}(x_{k+1})$, which project respectively to $p_k(u^-)$ and $p_k(x_{k+1})$ on V_k . As V_k is 8δ -quasiconvex, this lemma gives

$$d(p_k(u^-), p_k(x_{k+1})) \leq \max(21\delta, d(p_{k+1}(u^-), p_{k+1}(x_{k+1})) - d(p_{k+1}(u^-), p_k(u^-)) - d(p_{k+1}(x_{k+1}), p_k(x_{k+1})) + 42\delta).$$

As $d(p_k(u^-), p_k(x_{k+1})) \geq 22\delta$ by (2.10), the maximum has to be realized by the second term. Both distances $d(p_{k+1}(u^-), p_k(u^-))$ and $d(p_{k+1}(x_{k+1}), p_k(x_{k+1}))$ are equal to $2^k d^-$. We obtain

$$2 \cdot 2^k d^- - 20\delta \leq d(p_{k+1}(u^-), p_{k+1}(x_{k+1})).$$

As $d^- \geq D = 100\delta$, the left hand side is $\geq L - 8\delta = 92\delta$. This concludes the proof of (2.8), and of the induction.

Finally, if the conclusion of the lemma does not hold, then the induction will go on forever. Taking in particular $w = u^-$ in (2.9), we get $d(f(u^-), p_0(u^-)) \geq (2^{k+1} - 1)d^-$ for all k , a contradiction. \square

Here are some ways to optimize the proof to get better constants. In addition to multiple minor optimizations, let us mention the main ones:

- The set V_0 is 0-quasiconvex, not only 8δ -quasiconvex. This means that estimates in the proof of Lemma 2.3 are better for $k = 0$. There is a different source of gain for $k > 0$, thanks to the factor 2^k . Separating the two cases improves the final constant.
- There is an exponential gain in (2.12). One can spend some part of this gain, say $\exp(-(1 - \alpha)d' \log(2)/(10\delta)) \leq \exp(-(1 - \alpha)D \log(2)/(10\delta))$ to improve the multiplicative constant, and use the remaining part $\exp(-\alpha d' \log(2)/(10\delta))$ for the induction (for a suitable value of α).
- Instead of formulating the induction in terms of the distance from $f(z)$ to a geodesic G between $f(u^-)$ and $f(u^+)$, it is more efficient to induce over the Gromov product $(f(u^-), f(u^+))_{f(z)}$ (which coincides with the distance $d(f(z), G)$ up to 2δ) as most inequalities are done in terms of Gromov products. The main interest of this change is that, with the current argument, the point $f(u^-)$ projects on H between π_z and $f(z)$ within distance 4δ of π_z , which means there is a small loss. With the Gromov product approach, let m denote the point on G which is opposite to $f(z)$ in the triangle $[f(z), f(u^-), f(u^+)]$, i.e., it is on G at distance $(f(z), f(u^+))_{f(u^-)}$ of $f(u^-)$ and at distance $(f(z), f(u^-))_{f(u^+)}$ of $f(u^+)$. Let π_z denote the point on a geodesic H from $f(z)$ to m at distance $(f(u^-), f(u^+))_{f(z)}$ of $f(z)$. This point is within distance 2δ of m . It turns out that the projection of $f(u^-)$ on H is between m and π_z , i.e., opposite from $f(z)$. The above loss is suppressed in this approach.
- Finally, one can choose freely L , D and α within some range. In particular, L and D do not have to coincide. One can optimize numerically over these parameters to get the best possible bound. In the end, we take $L = 18\delta$ and $D = 55\delta$ and $\alpha = 12/100$ to get the value 92 in Theorem 1.1.

References

- [1] Mario Bonk, Oded Schramm, Embeddings of Gromov hyperbolic spaces, *Geom. Funct. Anal.* 10 (2000) 266–306. [MR1771428](#).
- [2] Martin R. Bridson, André Haefliger, *Metric Spaces of Non-Positive Curvature*, Grundlehren Math. Wiss. (Fundamental Principles of Mathematical Sciences), vol. 319, Springer-Verlag, Berlin, 1999. [MR1744486](#).
- [3] Michel Coornaert, Thomas Delzant, Athanase Papadopoulos, *Géométrie et théorie des groupes*, in: *Les groupes hyperboliques de Gromov (Gromov Hyperbolic Groups)*, in: *Lecture Notes in Math.*, vol. 1441, Springer-Verlag, Berlin, 1990, with an English summary. [1075994](#).
- [4] Sébastien Gouëzel, Gromov hyperbolicity, *Archive of Formal Proofs*, http://devel.isa-afp.org/entries/Gromov_Hyperbolicity.html, 2018, Formal proof development.
- [5] Vladimir Shchur, A quantitative version of the Morse lemma and quasi-isometries fixing the ideal boundary, *J. Funct. Anal.* 264 (3) (2013) 815–836. [3003738](#).
- [6] Vladimir Shchur, *Quasi-Isometries Between Hyperbolic Metric Spaces, Quantitative Aspects, Theses*, Université Paris Sud - Paris XI, July 2013.