


Provided for non-commercial research and educational use only.  
Not for reproduction or distribution or commercial use.



This article was originally published in a journal published by Elsevier in cooperation with Mendeleev Communications, and the attached copy is provided for the author's benefit and for the benefit of the author's institution, for non-commercial research and educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are prohibited. For exceptions, permission may be sought for such use through Elsevier's permissions site at:

<http://www.elsevier.com/locate/permissionusematerial>

## (Cyclopentadienyl)neodymium borohydrides with auxiliary N<sub>3</sub>-heterocyclic ligands

Guzelia I. Sadrtidinova,<sup>a,b</sup> Daniil A. Bardonov,<sup>a,b,c</sup> Konstantin A. Lyssenko,<sup>d</sup> Mikhail E. Minyaev,<sup>a,c</sup> Ilya E. Nifant'ev<sup>a,b,e</sup> and Dmitrii M. Roitershtein<sup>\*a,b,c</sup>

<sup>a</sup> A. V. Topchiev Institute of Petrochemical Synthesis, Russian Academy of Sciences, 119991 Moscow, Russian Federation. E-mail: roiter@yandex.ru

<sup>b</sup> National Research University Higher School of Economics (HSE University), 101000 Moscow, Russian Federation


<sup>c</sup> N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 119991 Moscow, Russian Federation

<sup>d</sup> G. V. Plekhanov Russian University of Economics, 117997 Moscow, Russian Federation

<sup>e</sup> Department of Chemistry, M. V. Lomonosov Moscow State University, 119991 Moscow, Russian Federation

DOI: 10.1016/j.mencom.2023.04.019

Mononuclear [CpNd(BH<sub>4</sub>)<sub>2</sub>L] borohydrides comprising L = 1,3,5-trimethyl-1,3,5-triazacyclohexane (Me<sub>3</sub>tach) and L = 1,4,7-trimethyl-1,4,7-triazacyclononane (Me<sub>3</sub>tacn) were obtained and structurally characterized. Their catalytic activity in coordination–insertion ring-opening polymerization of ε-caprolactone was studied.




**Keywords:** lanthanide complexes, neodymium complexes, cyclopentadienyl, borohydrides, crystal structure, polymerization, ε-caprolactone.

Cyclopentadienyl borohydride lanthanide complexes have great potential in the rare-earth chemistry due to the higher covalency of the metal (Ln)–ligand (BH<sub>4</sub>) bond, compared with the lanthanide halide analogs, since the borohydride ligand is more electron-donating.<sup>1–4</sup> Another attractive feature of borohydride complexes is their structural diversity due to different types of coordination of the borohydride ligand with the Ln<sup>3+</sup> cation, namely, tri-hapto (η<sup>3</sup>), di-hapto (η<sup>2</sup>), and even mono-hapto (η<sup>1</sup>) mode.<sup>5–10</sup> The latter easily becomes a drawback resulting in a mixture of products in the synthesis of borohydride complexes. We recently reported on the synthesis of a series of La, Nd, and Tb borohydride complexes.<sup>11</sup> A wide variety of structural types of cyclopentadienyl borohydride complexes of neodymium and lanthanum was observed, including five different structural types of Cp–lanthanide borohydrides. Three of them showed satisfactory activity in the polymerization reaction of ε-caprolactone. We assumed that in order to control the formation of a certain structural type in the Cp–Ln–BH<sub>4</sub> series, it is necessary to use an auxiliary ligand that effectively blocks the lanthanide ion coordination sphere. We have previously shown, using rare-earth arylcyclopentadienyl halide complexes, that saturated N-heterocyclic compounds such as 1,3,5-trimethyl-1,3,5-triazacyclohexane (Me<sub>3</sub>tach) and 1,4,7-trimethyl-1,4,7-triazacyclononane (Me<sub>3</sub>tacn) were highly effective in blocking the Ln<sup>3+</sup> coordination sphere.<sup>12</sup> Therefore, the goal of this work is the synthesis and characterization of cyclopentadienyl neodymium borohydrides containing Me<sub>3</sub>tach or Me<sub>3</sub>tacn ligands, and the study of their catalytic activity in the coordination–insertion ring-opening polymerization of ε-caprolactone.

Complexes [CpNd(BH<sub>4</sub>)<sub>2</sub>(Me<sub>3</sub>tach)] **1** and [CpNd(BH<sub>4</sub>)<sub>2</sub>(Me<sub>3</sub>tacn)] **2** were obtained by the reaction of


[CpNd(BH<sub>4</sub>)<sub>2</sub>(THF)<sub>2</sub>] with an excess of Me<sub>3</sub>tach or Me<sub>3</sub>tacn in THF (Scheme 1). Crystallization from THF/hexane solutions gave single crystals of **1** and **2**, respectively, suitable for X-ray diffraction. Both complexes are insoluble in nonpolar solvents and moderately soluble in THF, although markedly worse than their synthetic precursor [CpNd(BH<sub>4</sub>)<sub>2</sub>(THF)<sub>2</sub>]. Complex **2** is less soluble in THF than complex **1**; a similar trend in solubility was observed for the (triphenylcyclopentadienyl)neodymium chloride complexes.<sup>12</sup>

The structures of complexes **1** and **2** were established by X-ray diffraction analysis.<sup>†</sup> In both crystal structures under




Scheme 1

<sup>†</sup> Crystal data for **1**. C<sub>11</sub>H<sub>28</sub>B<sub>2</sub>N<sub>3</sub>Nd, *M* = 368.22, orthorhombic, space group *Pna*2<sub>1</sub>, *T* = 100 K, *a* = 15.2755(2), *b* = 7.80440(10), *c* = 13.5623(2) Å, *V* = 1616.84(4) Å<sup>3</sup>, and *Z* = 4. A total of 65492 (2θ<sub>max</sub> = 66.3°, *R*<sub>int</sub> = 0.0344)



**Figure 1** General view of complex **1**. The atoms are shown as thermal ellipsoids ( $p = 50\%$ ). Except for  $BH_4$ , all hydrogen atoms are omitted for clarity.



**Figure 2** General view of complex **2**. The atoms are shown as thermal ellipsoids ( $p = 50\%$ ), hydrogen atoms are omitted for clarity.

study, the  $Nd^{3+}$  cations are coordinated by a cyclopentadienyl ligand, three nitrogen atoms of the triaza ligand, and two borohydride ligands giving the neodymium cations a coordination number of eight (Figures 1, 2). The main geometrical parameters of complexes **1** and **2** are similar, despite significant differences in the size of the triaza ligands they contain. The  $Nd-Cp_{cent}$  distances are 2.489(13) Å (**1**) and 2.452(7) Å (**2**). The  $Nd-N$  distances are in the range of 2.642(2)–2.698(2) Å for complex **1** and 2.637(9)–2.655(12) Å for complex **2**. The major structural difference between **1** and **2** is the mutual arrangement of the cyclopentadienyl and the triaza ligands. Thus, the angle between the plane of the cyclopentadienyl ligand and the plane defined by the three nitrogen atoms is  $60.2^\circ$  for **1** and  $46.0^\circ$  for **2**. Apparently, such a noticeable difference is due to the fact that the bulkier  $Me_3tactn$  ligand in complex **2** experiences stronger repulsion from the cyclopentadienyl ligand than the  $Me_3tach$  ligand in complex **1**, which leads to a decrease in the angle between the corresponding planes. The consequence of this is a decrease in the coordination gap aperture.

The behavior of complexes **1** and **2** in solution is significantly different, despite their structural and chemical similarity. Thus, the  $^1H$  and  $^{11}B$  NMR spectra of complex **1** in  $THF-d_8$  indicate that the structure of **1** in solution is retained. At the same time, the  $^1H$  NMR spectrum of the analytically pure sample **2**, along with the signals of **2**, contains the signals for the free  $Me_3tactn$

reflections were collected and 6141 independent reflections were used for the structure solution and refinement, which converged to  $R_1 = 0.0143$  (for 5523 observed reflections),  $wR_2 = 0.0269$ , and  $GOF = 1.105$ .

*Crystal data for 2.*  $C_{14}H_{34}B_2N_3Nd$ ,  $M = 410.30$ , tetragonal, space group  $P4_3$ ,  $T = 120$  K,  $a = 9.754(2)$  Å,  $c = 19.962(6)$  Å,  $V = 1898.6(10)$  Å<sup>3</sup>, and  $Z = 4$  ( $Z' = 1$ ). A total of 16492 ( $2\theta_{max} = 58^\circ$ ,  $R_{int} = 0.0916$ ) reflections were collected and 5024 independent reflections were used for the structure solution and refinement. The refinement of the disordered part of the molecule was performed with EADP and DFIX instructions. Refinement converged to  $R_1 = 0.0554$  (for 4454 observed reflections),  $wR_2 = 0.1328$ , and  $GOF = 1.022$ .

CCDC 2221517 and 2222289 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk>.

ligand and other neodymium borohydride and triazacyclononane complexes (for the NMR spectra images, see Online Supplementary Materials, Figures S1–S5). Apparently, complex **2** is kinetically labile and undergoes redistribution reactions.

It is known that lanthanide borohydride complexes exhibit catalytic activity in the coordination–insertion ring-opening polymerization of  $\epsilon$ -caprolactone. The catalytic activity of such complexes depends significantly on the ligand environment of the lanthanide ion.<sup>13–19</sup> We have previously reported that mono- and bis(cyclopentadienyl)neodymium borohydrides could be used as such catalysts.<sup>11</sup> It was of undoubted interest to compare the catalytic activity of complexes **1** and **2** with the activity of those cyclopentadienyl neodymium borohydrides, especially knowing that in the polymerization of ethylene, the catalytic activity of triphenylcyclopentadienyl neodymium dichloride complexes would increase with the introduction of the  $Me_3tach$  ligand.<sup>12</sup>

To estimate the catalytic performance of **1** and **2** in comparison with the cyclopentadienyl neodymium borohydride complexes of a similar structure, but not containing N-donor ligands, namely  $[CpNd(BH_4)_2(THF)_2]$  **3** and  $[Cp_2Nd(BH_4)(THF)]$  **4**, polymerization was carried out under conditions similar to those described earlier (Table 1).<sup>11</sup> Both complexes exhibit higher activity in the polymerization of caprolactone compared with the activity of **3** and **4**, but only in the range of relatively high catalyst loadings, significantly decreasing at a catalyst/substrate molar ratio of 1:1000. Both complexes provide polymers with rather narrow polydispersity indices.

Interestingly, complex **2** shows higher activity than **1** despite the greater steric loading of **2** (containing the  $Me_3tactn$  ligand) compared to **1** (containing the  $Me_3tach$  ligand). There are at least two reasons for this. The first reason is that the initiation of the polymerization process begins with the interaction of the monomer molecules with the borohydride ligand, the availability of which depends only slightly on the coordination gap aperture. The second reason is the higher lability of  $Me_3tactn$  compared with that of  $Me_3tach$  in neodymium complexes. However, more research is needed to explain the advantage in using triazacyclononane ligand in the design of catalytic systems for caprolactone ring-opening polymerization, compared with triazacyclohexane.

In summary, auxiliary triaza heterocyclic ligands were used in the preparation of new cyclopentadienyl borohydride neodymium complexes possessing well-defined structures, in contrast to their precursors. The obtained complexes can effectively initiate  $\epsilon$ -caprolactone polymerization, surpassing their synthetic precursors in catalytic activity.

**Table 1** Polymerization of  $\epsilon$ -caprolactone (CL) initiated by complexes **1** and **2**.<sup>a</sup>

Entry	Initiator	CL/Nd ratio	1 min conversion <sup>b</sup> (%)	$t/min$	Yield <sup>c</sup> /g (%)	$M_{n,exp}^d / \times 10^{-3} g mol^{-1}$	$\bar{D}^d$
1	<b>1</b>	250	16	2.5	1.78 (89)	33.7	1.39
2	<b>1</b>	500	1	4.5	1.68 (84)	51.4	1.41
3	<b>1</b>	750	n.d.	6	1.15 (58)	61.7	1.17
4	<b>1</b>	1000	n.d.	>60	0.75 (38)	33.0	1.51
5	<b>2</b>	250	93	0.66	1.85 (93)	29.9	1.34
6	<b>2</b>	500	37	1	1.89 (94)	41.9	1.59
7	<b>2</b>	750	14	1.33	1.84 (92)	54.4	1.55
8	<b>2</b>	1000	5	>60	0.58 (29)	43.3	1.35

<sup>a</sup>General polymerization conditions: 1 M CL in THF and  $25^\circ C$ . <sup>b</sup>Monomer conversion in 1 min determined gravimetrically by the polymer yield in a separate series of experiments. <sup>c</sup>Yield: weight of the polymer obtained/weight of monomer, used after the reaction stopped. <sup>d</sup>Measured by GPC relative to polystyrene standards,  $\bar{D} = M_w/M_n$ .

The publication was prepared within the framework of the Academic Fund Program at HSE University in 2021–2022 (grant no. 21-04-034).

The authors are grateful for the exploitation of the equipment of the TIPS RAS Center of Collective Use: ‘New petrochemical processes, polymer composites and adhesives’.

#### Online Supplementary Materials

Supplementary data associated with this article can be found in the online version at doi: 10.1016/j.mencom.2023.04.019.

#### References

- 1 T. J. Marks and J. R. Kolb, *Chem. Rev.*, 1977, **77**, 263.
- 2 Z. Xu and Z. Lin, *Coord. Chem. Rev.*, 1996, **156**, 139.
- 3 M. Visseaux and F. Bonnet, *Coord. Chem. Rev.*, 2011, **255**, 374.
- 4 F. Ortu, *Chem. Rev.*, 2022, **122**, 6040.
- 5 V. D. Makhaev, *Russ. Chem. Rev.*, 2000, **69**, 727 (*Usp. Khim.*, 2000, **69**, 795).
- 6 M. Ephritikhine, *Chem. Rev.*, 1997, **97**, 2193.
- 7 F. Ortu, D. Packer, J. Liu, M. Burton, A. Formanuk and D. P. Mills, *J. Organomet. Chem.*, 2018, **857**, 45.
- 8 M. Visseaux, P. Zinck, M. Terrier, A. Mortreux and P. Roussel, *J. Alloys Compd.*, 2008, **451**, 352.
- 9 M. L. Khokhlov, A. E. Miroslavov, E. K. Legin, N. A. Korsakova, A. I. Kostylev, V. V. Gurzhiy, A. Yu. Ivanov and P. M. Tolstoi, *Mendeleev Commun.*, 2019, **29**, 696.
- 10 F. Bonnet, M. Visseaux, D. Barbier-Baudry, A. Hafid, E. Vigier and M. M. Kubicki, *Inorg. Chem.*, 2004, **43**, 3682.
- 11 D. A. Bardonov, P. D. Komarov, G. I. Sadrtidinova, V. K. Besprozvannyh, K. A. Lyssenko, A. O. Gudovannyi, I. E. Nifant'ev, M. E. Minyaev and D. M. Roitershtein, *Inorg. Chim. Acta*, 2022, **529**, 120638.
- 12 D. A. Bardonov, P. D. Komarov, V. I. Ovchinnikova, L. N. Puntus, M. E. Minyaev, I. E. Nifant'ev, K. A. Lyssenko, V. M. Korshunov, I. V. Taidakov and D. M. Roitershtein, *Organometallics*, 2021, **40**, 1235.
- 13 W. Li, M. Xue, Y. Zhang, Y. Yao and Q. Shen, *Z. Anorg. Allg. Chem.*, 2014, **640**, 1455.
- 14 S. M. Guillaume, M. Schappacher and A. Soum, *Macromolecules*, 2003, **36**, 54.
- 15 F. Yuan, T. Li, L. Li and Y. Zhou, *J. Rare Earths*, 2012, **30**, 753.
- 16 F. Jaroschik, F. Bonnet, X.-F. Le Goff, L. Ricard, F. Nief and M. Visseaux, *Dalton Trans.*, 2010, **39**, 6761.
- 17 I. Palard, A. Soum and S. M. Guillaume, *Macromolecules*, 2005, **38**, 6888.
- 18 D. Robert, M. Kondracka and J. Okuda, *Dalton Trans.*, 2008, **6**, 2667.
- 19 A. Momin, F. Bonnet, M. Visseaux, L. Maron, J. Takats, M. J. Ferguson, X. F. Le Goff and F. Nief, *Chem. Commun.*, 2011, **47**, 12203.

Received: 5th December 2022; Com. 22/7060